



# Trigonometry

## Graphs:


□  $y = \sin x$




□  $y = \cos x$




□  $y = \tan x$




□  $y = \cot x$



□  $y = \sec x$



□  $y = \csc x$



## Imp Results:

- $\sin(A+B)\sin(A-B) = \sin^2 A - \sin^2 B = \cos^2 B - \sin^2 A$
- $\cos(A+B)\cos(A-B) = \cos^2 A - \sin^2 B = \cos^2 B - \sin^2 A$
- $\sin(A+B+C) = \sin A \cos B \cos C + \cos A \sin B \cos C + \cos A \cos B \sin C - \sin A \sin B \sin C$
- $\cos(A+B+C) = \cos A \cos B \cos C - \cos A \sin B \sin C - \sin A \cos B \sin C - \sin A \sin B \cos C$
- $\tan(A+B+C) = \frac{\tan A + \tan B + \tan C - \tan A \tan B \tan C}{1 - \tan A \tan B - \tan B \tan C - \tan C \tan A}$

## Multiple angles

- $\sin 2A = 2 \sin A \cos A = \frac{2 \tan A}{1 + \tan^2 A}$
- $\cos 2A = \cos^2 A - \sin^2 A = 1 - 2 \sin^2 A = 2 \cos^2 A - 1 = \frac{1 - \tan^2 A}{1 + \tan^2 A}$
- $\tan 2A = \frac{2 \tan A}{1 - \tan^2 A}$

- $\sin 3A = 3 \sin A - 4 \sin^3 A$
- $\cos 3A = 4 \cos^3 A - 3 \cos A$
- $\tan 3A = \frac{3 \tan A - \tan^3 A}{1 - 3 \tan^2 A}$
- $\tan^2 A = \frac{1 - \cos 2A}{1 + \cos 2A}$
- $\cot 3A = \frac{\cot^3 A - 3 \cot A}{3 \cot^2 A - 1}$

## GP of Angles

$\cos A \cos 2A \cos 2^2 A \dots \cos 2^{n-1} A = \frac{\sin 2^n A}{2^n \sin A}$     □  $A+B+C = \frac{\pi}{2}$

## Inequalities: (AABC)

- $\tan A + \tan B + \tan C \geq 3\sqrt{3}$  [All acute]
- $\cos A + \cos B + \cos C \leq \frac{3}{2}$
- $\sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2} \leq \frac{1}{8}$
- $\sec A + \sec B + \sec C \geq 6$  [Acute]
- $\csc \frac{A}{2} + \csc \frac{B}{2} + \csc \frac{C}{2} \geq 6$  [Acute]

## Sum and Difference of two angles:

- $\sin(A+B) = \sin A \cos B + \cos A \sin B$
- $\sin(A-B) = \sin A \cos B - \cos A \sin B$
- $\cos(A+B) = \cos A \cos B - \sin A \sin B$
- $\cos(A-B) = \cos A \cos B + \sin A \sin B$
- $\tan(A \pm B) = \frac{\tan A \pm \tan B}{1 \mp \tan A \tan B}$
- $\cot(A \pm B) = \frac{\cot A \cot B \mp 1}{\cot A + \cot B}$

## Range of $f(\theta) = a \sin \theta + b \cos \theta$

$-\sqrt{a^2+b^2} \leq a \sin \theta + b \cos \theta \leq \sqrt{a^2+b^2}$

## Product into sum and difference

- $2 \sin A \sin B = \sin(A+B) - \sin(A-B)$
- $2 \cos A \sin B = \sin(A+B) + \sin(A-B)$
- $2 \cos A \cos B = \cos(A+B) + \cos(A-B)$
- $2 \sin A \cos B = \cos(A-B) - \cos(A+B)$

## Sum of difference into product

- $\sin C + \sin D = 2 \sin \left(\frac{C+D}{2}\right) \cos \left(\frac{C-D}{2}\right)$
- $\sin C - \sin D = 2 \cos \left(\frac{C+D}{2}\right) \sin \left(\frac{C-D}{2}\right)$
- $\cos C + \cos D = 2 \cos \left(\frac{C+D}{2}\right) \cos \left(\frac{C-D}{2}\right)$
- $\cos C - \cos D = 2 \sin \left(\frac{C+D}{2}\right) \sin \left(\frac{D-C}{2}\right)$

## Special Trios:

- $\cos A \cos(60-A) \cos(60+A) = \frac{\cos 3A}{4}$
- $\sin A \sin(60-A) \sin(60+A) = \frac{\sin 3A}{4}$
- $\tan A \tan(60-A) \tan(60+A) = \tan 3A$

## AP of Angles:

$\sin \alpha + \sin(\alpha+\beta) + \dots + \sin[\alpha+(n-1)\beta] = \frac{\sin \frac{n\beta}{2}}{\sin \frac{\beta}{2}} \left[ \sin \left( \alpha + \frac{(n-1)\beta}{2} \right) \right]$

## Conditional Identities ( $A+B+C = \pi$ )

- $\tan A + \tan B + \tan C = \tan A \tan B \tan C$
- $\tan \frac{A}{2} \tan \frac{B}{2} + \tan \frac{B}{2} \tan \frac{C}{2} + \tan \frac{C}{2} \tan \frac{A}{2} = 1$
- $\sin 2A + \sin 2B + \sin 2C = 4 \sin A \sin B \sin C$
- $\cos A + \cos B + \cos C = 1 + 4 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$
- $\sin A + \sin B + \sin C = 4 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}$

- $\sin^2 A + \sin^2 B + \sin^2 C = 1 - 2 \sin A \sin B \sin C$
- $\cos^2 A + \cos^2 B + \cos^2 C = 2 + 2 \sin A \sin B \sin C$
- In  $\Delta ABC$  □  $\cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} + \cos^2 \frac{C}{2} = 2 \cos \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}$
- $\cos^2 \frac{A}{2} + \cos^2 \frac{B}{2} + \cos^2 \frac{C}{2} = 2 + 2 \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$
- $\sin \frac{A}{2} + \sin \frac{B}{2} + \sin \frac{C}{2} \leq \frac{3}{2}$  □  $\tan^2 \frac{A}{2} + \tan^2 \frac{B}{2} + \tan^2 \frac{C}{2} \geq 1$

□  $\tan 3A - \tan 2A - \tan A = \tan 3A \tan 2A \tan A$

# Progression & Series

• Arithmetic Progression -  $a, a+d, a+2d, \dots, a+(n-1)d$  • nth term (General term):  $t_n = a+(n-1)d$ ,  $t_n = l-(n-1)d$

• 3 terms in AP consideration -  $a-d, a, a+d$  • 4 terms in A.P. -  $a-3d, a-d, a+d, a+3d$

• If  $a_1, a_2, a_3, \dots, a_{n-2}, a_{n-1}, a_n$  are in A.P.,  $a_1 + a_n = a_2 + a_{n-1} = a_3 + a_{n-2} = \dots = a_r + a_{n-r+1}$

• Sum of n terms in an AP:  $S_n = \frac{n}{2} [2a + (n-1)d] = \frac{n}{2} [a+l]$  • If  $S_n = an^2 + bn + c$ ,  $t_n = S_n - S_{n-1}$

• Arithmetic mean:  $AM(x_1, x_2) = \frac{x_1+x_2}{2}$ ,  $AM(x_1, x_2, x_3, \dots, x_n) = \frac{x_1+x_2+x_3+\dots+x_n}{n}$

•  $A_1 + A_2 + \dots + A_n = n \left( \frac{a+b}{2} \right)$

• Geometric Progression -  $a, ar, ar^2, \dots, a^{n-1}$  • nth term (General term):  $t_n = ar^{n-1}$

• Sum of n terms:  $S_n = \frac{a(1-r^n)}{1-r}$  [ $r > 1$ ] =  $\frac{a(r^n-1)}{r-1}$  [ $r > 1$ ] • If  $x_1, x_2, x_3, \dots$  are in GP,  $\log x_1, \log x_2, \dots$  are in AP

• 3 terms in GP:  $\frac{a}{r}, a, ar$  • 4 terms in GP:  $\frac{a}{r^3}, \frac{a}{r}, ar, ar^3$  • Sum of Infinite GP:  $S_\infty = \frac{a}{1-r}$  [ $|r| < 1$ ]

• Geometric means:  $GM(x_1, x_2) = (x_1 x_2)^{1/2}$ ,  $GM(x_1, x_2, x_3, \dots, x_n) = (x_1 x_2 x_3 \dots x_n)^{1/n}$

•  $G_1, G_2, G_3, \dots, G_n = (\sqrt[n]{ab})^n$

• Harmonic progression:  $a_1, a_2, a_3, \dots$  are in H.P. if  $\frac{1}{a_1}, \frac{1}{a_2}, \frac{1}{a_3}, \dots$  are in A.P.

• nth term (General term):  $t_n = \frac{1}{t_n \text{ of AP}}$  • Harmonic Mean:  $HM(x_1, x_2) = \frac{2x_1 x_2}{x_1 + x_2}$ ,  $HM(x_1, x_2, \dots, x_n) = \frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n}}$

• Inequalities:  $A \geq G \geq H$  •  $G^2 = AH$

• Special series:  $\sum n = \frac{n(n+1)}{2}$ ,  $\sum n^2 = \frac{n(n+1)(2n+1)}{6}$ ,  $\sum n^3 = \frac{n^2(n+1)^2}{4}$

• Arithmetic Geometric Progression (AGP):  $S = a + (a+d)r + (a+2d)r^2 + \dots$

$$S = a + (a+d)r + (a+2d)r^2 + \dots$$

$$-rS = \frac{ar}{r} + (a+d)r^2 + \dots$$

$$S(1-r) = a + d(r+r^2+\dots)$$

• Difference Series:  $S = 1 + 2 + 4 + 7 + 11 + 16 + \dots + t_n$   
 $S = 1 + 2 + 4 + 7 + 11 + \dots + t_n$   
 $0 = 1 + (1+2+3+\dots) - t_n$

$$\sum_{r=1}^n \frac{1}{r(r+1)(r+2)(r+3)} = \sum_{r=1}^n \frac{1}{3} \left[ \frac{(r+3)-r}{r(r+1)(r+2)(r+3)} \right] \quad \bullet \quad \sum_{r=1}^n r(r+1)(r+2)(r+3) = \sum_{r=1}^n \frac{1}{5} ((r+4) - (r-1)) (r(r+1)(r+2)(r+3))$$

Weighted mean:  $\frac{a_1^m + a_2^m + \dots + a_n^m}{n} > \left( \frac{a_1 + a_2 + a_3 + \dots + a_n}{n} \right)^m$  if  $m < 0$  or  $m > 1$

$\frac{a_1^m + a_2^m + \dots + a_n^m}{n} < \left( \frac{a_1 + a_2 + \dots + a_n}{n} \right)^m$  if  $0 < m < 1$



## Permutation & Combination

- Let  $p$  be a given prime and  $n$ , any positive integer, Then the maximum power of  $p$  present in  $n!$  is  $\left[\frac{n}{p}\right] + \left[\frac{n}{p^2}\right] + \left[\frac{n}{p^3}\right] + \dots$  where  $[\cdot]$   $\rightarrow$  Greatest Integer function
- Number of permutations of  $n$  different things taken  $r$  at a time  $\rightarrow {}^n P_r = \frac{n!}{(n-r)!}$
- Number of permutations of  $n$  different things taken all at a time  $\rightarrow n!$
- Number of permutation of  $n$  things [  $p$  are alike,  $q$  are alike,  $r$  are alike ]  $\rightarrow \frac{n!}{p!q!r!}$
- Number of combinations (selections) of  $n$  different things taking  $r$  at a time  $\rightarrow {}^n C_r = \frac{n!}{r!(n-r)!}$
- ${}^n C_r = {}^n C_{n-r}$ ,  ${}^n C_r + {}^n C_{r-1} = {}^{n+1} C_r$ ,  $r \cdot {}^n C_r = n \cdot {}^{n-1} C_{r-1}$ ,  $\frac{{}^n C_r}{r+1} = \frac{{}^{n+1} C_{r+1}}{n+1}$ ,  $\frac{{}^n C_r}{{}^n C_{r-1}} = \frac{n-r+1}{r}$
- When  $n$  is even, max value of  ${}^n C_r \rightarrow {}^n C_{n/2}$
- When  $n$  is odd, max value of  ${}^n C_r \rightarrow {}^n C_{\frac{n-1}{2}}$  or  ${}^n C_{\frac{n+1}{2}}$
- When ACW/CW doesn't matter (e.g. necklace, garland),  $\rightarrow$  circular arrangement  $\rightarrow \frac{(n-1)!}{2}$
- Total no of combination of  $n$  things taken 1 or more at a time  $\rightarrow {}^n C_1 + {}^n C_2 + {}^n C_3 + \dots + {}^n C_n = 2^n - 1$
- Total no of selections of  $n$  things, [  $p$  similar,  $q$  similar,  $r$  alike ]  $\rightarrow$  (including  $\emptyset$ )  $\rightarrow (p+1)(q+1)(r+1)$
- If  $N = p_1^a \times p_2^b \times p_3^c \times \dots$  where  $a, b, c, \dots$  are non-negative integers,  $p_1, p_2, p_3, \dots$  are prime no. Then  $\rightarrow$  Total No of Divisors =  $(a+1)(b+1)(c+1)\dots$  Sum of all divisors =  $\left(\frac{p_1^{a+1}-1}{p_1-1}\right) \times \left(\frac{p_2^{b+1}-1}{p_2-1}\right) \times \left(\frac{p_3^{c+1}-1}{p_3-1}\right) \times \dots$
- All the divisors excluding 1 and  $N$  are called proper divisors
- No of ways of writing  $N$  as a product of two natural nos  $\rightarrow \left\{ \begin{array}{l} \left[ \frac{1}{2} (a+1)(b+1)(c+1)\dots \right]$  if  $N$  isn't a perfect square \\  $\left[ \frac{1}{2} (a+1)(b+1)(c+1)\dots + 1 \right]$  if  $N$  is a perfect square \end{array} \right.
- $N$  is a perfect square if  $a, b, c, \dots$  all are even
- $N$  is a perfect cube if  $a, b, c, \dots$  all are multiples of 3.
- $N = 2^a \times 3^b \times 5^c \times \dots$  If  $N$  is odd,  $a=0, b, c, d, \dots \geq 0$  If  $N$  is even,  $a \geq 1, b, c, \dots \geq 0$
- No of Non negative integral sol<sup>n</sup> of the eq<sup>n</sup>  $x_1 + x_2 + x_3 + \dots + x_r = n$  is  $\rightarrow {}^{n+r-1} C_{r-1}$
- No of positive integral sol<sup>n</sup> of the eq<sup>n</sup>  $x_1 + x_2 + x_3 + \dots + x_r = n$  is  $\rightarrow {}^{n-1} C_{r-1}$
- Sum of all  $n$ -digit numbers formed using  $n$  digits =  $(n-1)! (\text{Sum of all } n \text{ digits}) \times (111\dots 1)_{n \text{ times}}$
- No of diagonals of  $n$  sided polygon  $\rightarrow \frac{n(n-3)}{2}$
- No of squares in two system of perpendicular parallel lines (when 1st set contain  $m$  lines and 2nd set contain  $n$  lines) is equal to  $\rightarrow \sum_{r=1}^{m-1} (m-r)(n-r)$ ; ( $m < n$ )
- Derangements: No of ways so that no letter goes to the correct address.  


$$D_n = n! \left[ 1 - \frac{1}{1!} + \frac{1}{2!} - \frac{1}{3!} + \frac{1}{4!} - \dots + (-1)^n \frac{1}{n!} \right]$$

# Complex Number

- $z = x + iy$ ,  $x, y \in \mathbb{R}$  and  $i = \sqrt{-1}$      $\text{Re}(z) = x$ ,  $\text{Im}(z) = y$      $\sqrt{-a} = i\sqrt{a}$
- The property  $\sqrt{a}\sqrt{b} = \sqrt{ab}$  is valid only if at least one of  $a$  and  $b$  is non negative, if  $a$  and  $b$  are both negative, then  $\sqrt{a}\sqrt{b} = -\sqrt{|a||b|}$
- $a+ib > c+id$  is meaningful only if  $b=d=0$     • If  $a+ib = c+id$ ,  $a=c$ ,  $b=d$
- In real no system,  $a^2+b^2=0$ ,  $a=b=0$ . But  $z_1^2+z_2^2=0$  does NOT mean  $z_1=z_2=0$ .
- $i = \sqrt{-1}$ ,  $i^2 = -1$ ,  $i^3 = -i$ ,  $i^4 = 1$ ;  $i^{4n+1} = i$ ,  $i^{4n+2} = -1$ ,  $i^{4n+3} = -i$ ,  $i^{4n} = 1$

Square Root of a Complex No:  $\sqrt{a+ib} = x+iy \Rightarrow a = x^2 - y^2$ ;  $2xy = b$  solve.  
Sign of  $b$  decides whether  $x$  and  $y$  are of same sign or opposite sign.





Modulus of CN:  $|z| = r = \sqrt{x^2+y^2}$



Amplitude of CN:

Argument/amplitude of CN,  $\theta = \tan^{-1}(y/x) = \tan^{-1}(\frac{\text{Im}(z)}{\text{Re}(z)})$   
From the real axis,  $\arg(z) \in [-\pi, \pi]$


Principle Argument



Polar form

•  $z = x + iy = r(\cos\theta + i\sin\theta)$

Euler's form  $z = r(\cos\theta + i\sin\theta) = r e^{i\theta}$   
 $\therefore e^{i\theta} = \cos\theta + i\sin\theta$



Properties of Conjugates


- $\overline{\bar{z}} = z$     • If  $z = \bar{z}$ ,  $z$  is purely real
- $z + \bar{z} = 0$ ,  $z$  is purely imaginary
- $z \cdot \bar{z} = z^2$     •  $\overline{z_1 + z_2} = \bar{z}_1 + \bar{z}_2$
- $\overline{z_1 z_2} = \bar{z}_1 \cdot \bar{z}_2$
- $\overline{z^n} = (\bar{z})^n$     •  $\overline{\left(\frac{z_1}{z_2}\right)} = \frac{\bar{z}_1}{\bar{z}_2}$

$z + \bar{z} = 2 \text{Re}(z)$   
 $z - \bar{z} = 2 \text{Im}(z)$

Properties of Arguments

- $\arg(z_1 z_2) = \arg(z_1) + \arg(z_2)$
- $\arg(z_1 z_2 \dots z_n) = \arg(z_1) + \dots + \arg(z_n)$
- $\arg\left(\frac{z_1}{z_2}\right) = \arg(z_1) - \arg(z_2)$
- $\arg(\bar{z}) = -\arg(z)$
- $\arg(z^n) = n \arg(z)$
- $\arg\left(\frac{1}{z}\right) = -\arg(z)$
- If  $z$  is purely imaginary,  $\arg(z) = \pm \pi/2$
- If  $z$  is purely real,  $\arg(z) = 0/\pi$


Angle b/w line joining  $z_1$  and  $z_2$  &  $z_3, z_4$



Properties of modulus

- $|z| = 0 \Rightarrow z = 0 = \text{Re}(z) = \text{Im}(z)$
- $|z| = |\bar{z}| = |-z| = |- \bar{z}|$
- $-|z| \leq \text{Re}(z) \leq |z|$
- $-|z| \leq \text{Im}(z) \leq |z|$
- $z \cdot \bar{z} = |z|^2$     •  $|z^n| = |z|^n$
- $|z_1 z_2 \dots z_n| = |z_1| |z_2| \dots |z_n|$
- $\left|\frac{z_1}{z_2}\right| = \frac{|z_1|}{|z_2|}$
- $|z_1 + z_2|^2 = |z_1|^2 + |z_2|^2 + 2\text{Re}(z_1 \bar{z}_2)$
- $|z_1 - z_2|^2 = |z_1|^2 + |z_2|^2 - 2\text{Re}(z_1 \bar{z}_2)$
- $|z_1 + z_2|^2 + |z_1 - z_2|^2 = 2(|z_1|^2 + |z_2|^2)$
- $|z_1 - z_2| \rightarrow$  dist b/w  $z_1$  &  $z_2$
- $|z_1 + z_2| \leq |z_1| + |z_2|$
- $|z_1 + z_2 + \dots + z_n| \leq |z_1| + |z_2| + \dots + |z_n|$
- $|z_1 + z_2| \geq ||z_1| - |z_2||$

Angle b/w 2 lines



• If  $z_1, z_2$  and  $z_3$  are vertices of an equilateral triangle. Then

$\frac{1}{z_1 - z_2} + \frac{1}{z_2 - z_3} + \frac{1}{z_3 - z_1} = 0$

i.e.  $z_1^2 + z_2^2 + z_3^2 = z_1 z_2 + z_2 z_3 + z_3 z_1$

• Square root of  $z = a + ib$  are

$$\left\{ \begin{array}{l} \pm \sqrt{\frac{|z|+a}{2}} + i \sqrt{\frac{|z|-a}{2}}, \text{ for } b > 0 \\ \pm \sqrt{\frac{|z|+a}{2}} - i \sqrt{\frac{|z|-a}{2}}, \text{ for } b < 0 \end{array} \right.$$



• If  $z_1, z_2, z_3$  are vertices of an isosceles right angled triangle, w/ right angle at  $z_3$ , then  
 $(z_1 - z_2)^2 = 2(z_1 - z_2)(z_1 - z_3)$

**De Moivre's Theorem**

$(\cos \theta + i \sin \theta)^n = (e^{i\theta})^n = e^{in\theta} = (\cos(n\theta) + i \sin(n\theta))$


- $(\cos \theta - i \sin \theta)^n = \cos n\theta - i \sin n\theta$     •  $\frac{1}{\cos \theta + i \sin \theta} = (\cos \theta - i \sin \theta)^{-1} = \cos \theta - i \sin \theta$
- $(\sin \theta \pm i \cos \theta)^n \neq \sin n\theta \pm i \cos n\theta$     •  $(\cos \theta, -i \sin \theta)^n \neq \cos n\theta, + i \sin n\theta$
- $(\sin \theta + i \cos \theta)^n = [\cos(\frac{\pi}{2} - \theta) + i \sin(\frac{\pi}{2} - \theta)]^n = [\cos n(\frac{\pi}{2} - \theta) + i \sin n(\frac{\pi}{2} - \theta)]$

**Cube Roots of Unity**

$z = 1^{\frac{1}{3}} = 1, \omega, \omega^2$  where  $\omega = -\frac{1}{2} + i\frac{\sqrt{3}}{2}, \omega^2 = -\frac{1}{2} - i\frac{\sqrt{3}}{2}$   
 $\omega = e^{i\frac{2\pi}{3}}, \omega^2 = e^{-i\frac{2\pi}{3}}$

- Sum of roots is 0;  $1 + \omega + \omega^2 = 0$     • Product of roots = 1;  $1 \cdot \omega \cdot \omega^2 = 1$
- $\omega = \frac{1}{\omega^2}, \omega^2 = \frac{1}{\omega}$     •  $\omega = \overline{\omega^2}, \omega^2 = \overline{\omega}$     •  $\omega^{3n-1} = \omega, \omega^{3n-2} = \omega^2, \omega^{3n} = 1$
- $1 + \omega^3 + \omega^{2n} = \begin{cases} 3, & n \text{ is a multiple of } 3 \\ 0, & n \text{ is not a multiple of } 3 \end{cases}$     • Cube roots of unity represent the vertices of an equilateral triangle on Argand Plane

**Section Formula**



- Centroid of  $\Delta$  formed by  $z_1, z_2$  and  $z_3 \rightarrow \frac{z_1 + z_2 + z_3}{3}$
- If circumcentre of an  $\Delta$  is origin, then orthocentre  $\rightarrow z_1 + z_2 + z_3$

**nth root of unity**

- Sum of all nth roots of unity = 0
- Product of all roots =  $1 \cdot \alpha \cdot \alpha^2 \cdot \alpha^3 \dots \alpha^{n-1} = \begin{cases} 1, & n \text{ is odd} \\ -1, & n \text{ is even} \end{cases}$

$z = 1^{\frac{1}{n}} \Rightarrow z = (\cos 2k\pi + i \sin 2k\pi)^{\frac{1}{n}} = \cos \frac{2k\pi}{n} + i \sin \frac{2k\pi}{n}$   
 $\therefore \alpha = \cos \frac{2\pi}{n} + i \sin \frac{2\pi}{n}$

**Locus of a CN** ( $z_1$  and  $z_2$  are fixed,  $z$  is a variable point)

$|z - z_1| = |z - z_2| \rightarrow z$  lies on perpendicular bisector of  $z_1z_2$

$|z - z_1| + |z - z_2| = |z_1 - z_2| \rightarrow z$  lies on the segment joining  $z_1$  and  $z_2$


$|z - z_1|^2 + |z - z_2|^2 = |z_1 - z_2|^2 \rightarrow$  Circle with  $z_1$  and  $z_2$  as diameter extremities.

$\arg\left(\frac{z - z_1}{z - z_2}\right) = \pm \frac{\pi}{2} \rightarrow \arg\left(\frac{z - z_1}{z - z_2}\right) = \frac{\pi}{2}$

$\arg\left(\frac{z - z_1}{z - z_2}\right) = \alpha$  (fixed)

# Straight Line

## Rotation of Axes



$$\begin{aligned} x &= x' \cos \theta - y' \sin \theta & x' &= x \cos \theta + y \sin \theta \\ y &= x' \sin \theta + y' \cos \theta & y' &= -x \sin \theta + y \cos \theta \end{aligned}$$

	x ↓	y ↓
x' ↓	cos θ	sin θ
y' ↓	-sin θ	cos θ

## Distance formula.

$$|d| = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2}$$

## Area of triangle

$$\Delta = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

## Shoebat Method

$$\Delta = \frac{1}{2} \begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \\ x_3 & y_3 \end{vmatrix} = \frac{1}{2} [(x_1 y_2 + x_2 y_3 + x_3 y_1) - (x_2 y_1 + x_3 y_2 + x_1 y_3)]$$

## Area of polygon

$$\Delta = \frac{1}{2} \begin{vmatrix} x_1 & y_1 \\ x_2 & y_2 \\ \vdots & \vdots \\ x_n & y_n \\ x_1 & y_1 \end{vmatrix}$$

Points must be taken in cyclic order.

## Section formula

Internal:  $(\frac{mx_2 + nx_1}{m+n}, \frac{my_2 + ny_1}{m+n})$

## Special points:

Centroid:  $(\frac{\Sigma x}{3}, \frac{\Sigma y}{3})$

Circumcentre:  $(\frac{x_1 \sin 2A + x_2 \sin 2B + x_3 \sin 2C}{\sin 2A + \sin 2B + \sin 2C}, \text{similar})$


External:  $(\frac{mx_2 - nx_1}{m-n}, \frac{my_2 - ny_1}{m-n})$

Incentre:  $(\frac{ax_1 + bx_2 + cx_3}{a+b+c}, \frac{ay_1 + by_2 + cy_3}{a+b+c})$

O, G, H of an acute angle triangle are collinear

$$O : G : H = 1 : 2$$


## Straight line



Eqn of line  $\rightarrow y = mx + c$

$$\frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$$

## Angle b/w two lines



$$\tan \theta = \pm \frac{m_1 - m_2}{1 + m_1 m_2}$$

If  $m_1 = m_2 \rightarrow$  lines are parallel

If  $m_1 m_2 = -1 \rightarrow$  lines are perpendicular.

## Eqn of line

- Parallel to x axis  $\rightarrow y = b$
- Parallel to y axis  $\rightarrow x = a$
- Normal form  $\rightarrow x \cos \alpha + y \sin \alpha = p$
- Slope Intercept form  $\rightarrow y = mx + c$
- Point slope form  $\rightarrow y - y_1 = m(x - x_1)$
- Intercept form  $\rightarrow \frac{x}{a} + \frac{y}{b} = 1$

Two point form  $\rightarrow \frac{y - y_1}{y_2 - y_1} = \frac{x - x_1}{x_2 - x_1}$

## Conditions

- Coincident if  $\frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$
- Parallel if  $\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}$
- Intersecting if  $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$
- Perpendicular if  $a_1 a_2 + b_1 b_2 = 0$

## Parametric form

$A(x, y) = (x_1 + r \cos \theta, y_1 + r \sin \theta)$

$ax + by + c = 0$

- Parallel to line  $\rightarrow ax + by + \lambda = 0$
- Perpendicular to line  $\rightarrow bx - ay + \mu = 0$

## Concurrence of three lines

$$\begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix} = 0$$

## Dist of a point from a line

$$|d| = \frac{|ax_1 + by_1 + c|}{\sqrt{a^2 + b^2}}$$

## Dist b/w two parallel lines

$$d = \frac{|c_1 - c_2|}{\sqrt{a^2 + b^2}}$$

## Angle bisector

of  $a_1 x + b_1 y + c_1 = 0$  &  $a_2 x + b_2 y + c_2 = 0$


$$\frac{a_1 x + b_1 y + c_1}{\sqrt{a_1^2 + b_1^2}} = \pm \frac{a_2 x + b_2 y + c_2}{\sqrt{a_2^2 + b_2^2}} \rightarrow (X)$$

$$\frac{a_1 x + b_1 y + c_1}{\sqrt{a_1^2 + b_1^2}} = - \frac{a_2 x + b_2 y + c_2}{\sqrt{a_2^2 + b_2^2}} \rightarrow (Y)$$

- Make  $c_1, c_2$  +ve
- Then (X) contains origin

	Acute bisec	Obtuse bisec	
$a_1 a_2 + b_1 b_2 > 0$	(Y)	(X)	Origin is obtuse
$a_1 a_2 + b_1 b_2 < 0$	(X)	(Y)	Origin is in acute.

## Family of St. lines



## For foot of Perpendicular

Image

A(x, y) B(x2, y2) C(x3, y3)

B/C  $\frac{x_2 - x_1}{a} = \frac{y_2 - y_1}{b} = - \frac{(ax_1 + by_1 + c)}{a^2 + b^2}$

C/B  $\frac{x_3 - x_1}{a} = \frac{y_3 - y_1}{b} = - \frac{2(ax_1 + by_1 + c)}{a^2 + b^2}$

## Pair of St. lines

$$(a_1 x + b_1 y + c_1)(a_2 x + b_2 y + c_2) = 0$$

## Bisector of Angle b/w pair of st. line

$$ax^2 + 2hxy + by^2 = 0$$

$$\frac{x^2 - y^2}{a - b} = \frac{xy}{h}$$

## Angle b/w pair of st. lines

$$m_1 + m_2 = -\frac{2h}{b}; m_1 m_2 = \frac{a}{b} \therefore \tan \theta = \left| \frac{2\sqrt{h^2 - ab}}{a + b} \right|$$

- $a + b = 0 \rightarrow$  lines are perpendicular
- $h^2 = ab \rightarrow$  lines are || or coincident.

## General 2nd degree equation

$$ax^2 + 2hxy + bx^2 + 2gx + 2fy + c = 0$$

Pair of st. line if

$$\Delta(abc + 2fgh - af^2 - bg^2 - ch^2) = 0$$

Point of intersection  $\rightarrow \left( \frac{bg - hf}{h^2 - ab}, \frac{af - gf}{h^2 - ab} \right)$

$\Delta \neq 0, h^2 > ab$   
(Hyperbola)

Ellipse  
( $\Delta \neq 0, h^2 < ab$ )

- $\Delta = 0$   
(Pair of st. line)
- $\Delta \neq 0, a = b, h = 0$   
(Circle)
- $\Delta \neq 0, h^2 = ab$   
(Parabola)



# Circle

## Eqn of Circle

• If centre  $(0,0) \rightarrow x^2 + y^2 = r^2$

• From diameter extremities:

$$(x-x_1)(x-x_2) + (y-y_1)(y-y_2) = 0$$

• Intercepts made on axis

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

x int  $\rightarrow 2\sqrt{g^2 - c}$   
y int  $\rightarrow 2\sqrt{f^2 - c}$

• Equation of circumcircle of  $\Delta$  formed by  $a_1x + b_1y + c_1 = 0$  w/ coordinate axis.

$$ab(x^2 + y^2) + c(bx + ay) = 0$$

## Tangents

$$x^2 + y^2 + 2gx + 2fy + c = 0$$

$$T \rightarrow xx_1 + yy_1 + g(x+x_1) + f(y+y_1) + c = 0$$

• For  $x^2 + y^2 = a^2$ :

Point form  $\rightarrow xx_1 + yy_1 - a^2 = 0$

Parametric  $\rightarrow x \cos \theta + y \sin \theta - a = 0$

Slope form  $\rightarrow y = mx \pm a\sqrt{1+m^2}$

• For  $x^2 + y^2 + 2gx + 2fy + c = 0$

$$y + f = m(x + g) \pm \sqrt{g^2 + f^2 - c} \sqrt{1+m^2}$$

• From a point outside

$$(y - y_1) = m(x - x_1)$$

• Length of tangent from a point to a circle

$$d = \sqrt{S_1}$$

• Pair of tangents (combined eqn)

$$SS_1 = T^2$$


## Length of common tangents

DCT  $\rightarrow |AB| = \sqrt{d^2 - (r_1 - r_2)^2}$

TCT  $\rightarrow |CD| = \sqrt{d^2 - (r_1 + r_2)^2}$

[d  $\rightarrow$  dist b/w two centres]

## Radical Centre



Solve,  
 $S_1 - S_2 = 0$   
 $S_2 - S_3 = 0$   
 $S_3 - S_1 = 0$

• Circles touch a line at  $(x_1, y_1)$ !






$$(x-x_1)^2 + (y-y_1)^2 + \lambda L = 0$$

• From general equation  $x^2 + y^2 + 2gx + 2fy + c = 0$   
 • Centre  $(-g, -f)$   
 • radius  $= \sqrt{g^2 + f^2 - c}$

• Two lines  $a_1x + b_1y + c_1 = 0$ ;  $a_2x + b_2y + c_2 = 0$  cut the coordinate axes at concyclic points. if  $m_1 m_2 = 1 \rightarrow a_1 a_2 = b_1 b_2$

• Parametric form  $x^2 + y^2 = r^2 \rightarrow (r \cos \theta, r \sin \theta) [0 \leq \theta < 2\pi]$   
 $(x-h)^2 + (y-k)^2 = r^2 \rightarrow (h + r \cos \theta, k + r \sin \theta)$

• A line intersects, touches or doesn't intersect the circle if radius is greater than, equal to or less than the length of perpendicular from centre of the circle to the line.



• If two lines  $a_1x + b_1y + c_1 = 0$  and  $a_2x + b_2y + c_2 = 0$  meet the axes in four distinct points concyclic points, then  $a_1 a_2 = b_1 b_2$  and also the eqn of the circle passing thru those concyclic points is:  
 $(a_1x + b_1y + c_1)(a_2x + b_2y + c_2) - (a_1 b_2 + a_2 b_1)xy = 0$

## Tangents

• Normals  $(y - y_1) = \frac{y_1 + f}{x_1 + g}(x - x_1)$

## Director circle

• Angle of Intersection of two circles:  
 • When  $\theta = 90^\circ$  [Orthogonally]  $\rightarrow 2g_1 g_2 + 2f_1 f_2 = c_1 + c_2$

• Chord of Contact  $\rightarrow T = 0$   
 • Eqn of chord bisected at  $(x_2, y_2) \rightarrow S_1 = T$

$$\cos \theta = \frac{r_1 r_2 - \sqrt{r_1^2 + r_2^2 - d^2}}{2r_1 r_2}$$

## Intersections

•  $|x_1 x_2| > r_1 + r_2$   
 [2 Direct  $T_s$ ]  
 [2 Transverse  $T_s$ ]  
 [P divided  $x_1, x_2$  externally in ratio  $r_1 : r_2$ ]


•  $|x_1 x_2| = r_1 + r_2$   
 [Total 3  $T_s$ ]

•  $|r_1 - r_2| < |x_1 x_2| < |r_1 + r_2|$   
 [Total 2 Common  $T_s$ ]


•  $|x_1 x_2| = |r_1 - r_2|$   
 [1 CT]

•  $|x_1 x_2| < |r_1 - r_2|$   
 [No CT]

## Common Chord






## Radical Axis




• RA is  $\perp$  to line joining the centres  
 • RA bisects common tangents  
 • Need not pass thru mp of  $x_1, x_2$   
 • If 2 circles cut a third circle orthogonally, RA of those 2 pass thru 3rd one's centre.

## Family of Circles



# PARABOLA

## Eccentricity




Equation of Conic:  
Focus  $(\alpha, \beta)$ , Directrix  $(ax + by + c = 0)$   
 $\rightarrow (x - \alpha)^2 + (y - \beta)^2 = e^2 \left( \frac{ax + by + c}{a^2 + b^2} \right)^2$

- $e = 0 \rightarrow$  circle
- $e = 1 \rightarrow$  Parabola
- $e < 1 \rightarrow$  Ellipse
- $e > 1 \rightarrow$  Hyperbola
- $e = \infty \rightarrow$  Pair of st. lines

Standard forms  $\rightarrow$   $y^2 = 4ax$   $\rightarrow$   $y^2 = -4ax$   $\rightarrow$   $x^2 = 4ay$   $\rightarrow$   $x^2 = -4ay$

Position of a point w.r.t. a Parabola  $y^2 = 4ax \rightarrow$   
 $y^2 - 4ax > 0 \rightarrow$  outside /  $y^2 - 4ax = 0 \rightarrow$  on /  $y^2 - 4ax < 0 \rightarrow$  Inside



$x = Ay^2 + By + C$   
Vertex  $\rightarrow (-\frac{B}{2A}, -\frac{D}{2A})$   
Length LR  $\rightarrow \frac{1}{|A|}$

## Parametric form

$(y - k)^2 = 4a(x - h)$   
 $\rightarrow x = h + at^2$   
 $y = k + 2at$

## Properties of Focal Chord

- Chord joining  $P(at_1^2, 2at_1)$  and  $Q(at_2^2, 2at_2)$  is a focal chord, then  $t_1 t_2 = -1$ ,  $Q \equiv (\frac{a}{t_1^2}, -\frac{2a}{t_1})$
- Focal chord from  $P(at^2, 2at)$  has length  $a(t + \frac{1}{t})^2$
- Focal chord making angle  $\theta$  with axis has length  $4a \operatorname{cosec}^2 \theta$
- Semi latus Rectum is HPM of SP & SQ where P and Q are extremities of focal chord, S  $\rightarrow$  focus
- Circle described on the focal length as diameter touches tangent at vertex
- Circle described on the focal chord as diameter touches directrix.
- Pair of tangents:  $\rightarrow SS_1 = T^2$   $\rightarrow T = 0$   $\rightarrow S_1 = T$

## Equation of tangents

- Point form  $\rightarrow yy_1 = 2a(x + x_1)$
- Parametric form  $\rightarrow ty = x + at^2$  [at  $(at^2, 2at)$ ]
- Slope form  $\rightarrow y = mx + \frac{a}{m}$  [at  $(\frac{a}{m^2}, \frac{2a}{m})$ ]

## Properties of Tangents

- Point of Intersection of tangents at two points  $A(at_1^2, 2at_1)$  and  $B(at_2^2, 2at_2)$  on the Parabola  $y^2 = 4ax$  is  $T(at_1 t_2, a(t_1 + t_2))$
- Locus of foot of  $\perp$  from focus upon any tangent is tangent at vertex.
- Length of Tangent b/w POC and POI w/ directrix subtends  $90^\circ$  at the focus.
- Tangents at extremities of focal chord are perpendicular on Directrix.



## Equation of Normal

- Point form  $\rightarrow y - y_1 = -\frac{y_1}{2a}(x - x_1)$
- Parametric form  $\rightarrow y = -tx + 2at + at^3$
- Slope form  $\rightarrow y = mx - 2am - am^3$

Parabola	Normal (Parametric)	Normal (Slope)
$y^2 = 4ax$	$y = -tx + 2at + at^3$ ( $at^2, 2at$ )	$y = mx - 2am - am^3$ ( $am^2, -2am$ )
$y^2 = -4ax$	$y = tx + 2at + at^3$ ( $-at^2, 2at$ )	$y = mx + 2am + am^3$ ( $-am^2, 2am$ )
$x^2 = 4ay$	$x = -ty + 2at + at^3$ ( $2at, at^2$ )	$y = mx + 2a + \frac{a}{m^2}$ ( $-\frac{2a}{m}, \frac{a}{m^2}$ )
$x^2 = -4ay$	$x = ty + 2at + at^3$ ( $2at, -at^2$ )	$y = mx - 2a - \frac{a}{m^2}$ ( $\frac{2a}{m}, -\frac{a}{m^2}$ )

## Properties of Normals


- Normals other than axis of Parabola never passes thru focus.
- POI of Normals from  $P(at_1^2, 2at_1)$ ,  $Q(at_2^2, 2at_2)$   $\rightarrow [2a + a(t_1^2 t_2^2 + t_1 t_2), -at_1 t_2 (t_1 + t_2)]$
- Normal at point  $P(t_1)$  meets the curve again at  $Q(t_2)$ ,  $t_2 = -t_1 - \frac{2}{t_1}$

## Co-normal Points

- $y = mx - 2am - am^3 \Rightarrow am^3 + (2a - h)m + k = 0$  (cubic in m)
- $m_1 + m_2 + m_3 = 0$  ;  $m_1 m_2 + m_2 m_3 + m_3 m_1 = \frac{2a - h}{a}$
- Algebraic sum of ordinates of co-normal points = 0
  - Centroid of the triangle formed by them lies on axis.
  - If three normals drawn on  $y^2 = 4ax$  from  $(h, k)$  is real  $\rightarrow |h| > 2a$

## Reflection Property of Parabola

The tangent at any point P to a parabola bisects the angle between the focal chord through P and perpendicular from P to the directrix




- Thus, if any light ray is sent along a line parallel to the axis of the parabola then the reflected ray passes thru the focus, as the normal bisects the angle between the incident ray and reflected ray.
- Tangents are drawn from the point  $(x_1, y_1)$  to the parabola  $y^2 = 4ax$ , the length of the chord of contact  $= \frac{1}{|a|} \sqrt{(y_1^2 - 4ax_1)(y_1^2 + 4ax_1)}$

- Area of the triangle formed by the tangents drawn from  $(x_1, y_1)$  to  $y^2 = 4ax$  and their chord of contact is  $\frac{(y_1^2 - 4ax_1)^{3/2}}{2a}$



# Properties of Triangle / Solution of Triangle



$a+b+c = 2s$  (Perimeter)

$s = \frac{a+b+c}{2}$  (semi perimeter)

Sine Rule

$\frac{\sin A}{a} = \frac{\sin B}{b} = \frac{\sin C}{c} = 2R$   
[R → circumradius]


Cosine Rule

#  $\cos A = \frac{b^2+c^2-a^2}{2bc}$

#  $\cos B = \frac{c^2+a^2-b^2}{2ca}$

#  $\cos C = \frac{a^2+b^2-c^2}{2ab}$

Projection Formula



#  $a = b \cos C + c \cos B$

#  $b = c \cos A + a \cos C$

#  $c = a \cos B + b \cos A$

Napier Formula

#  $\tan\left(\frac{A-B}{2}\right) = \frac{a-b}{a+b} \cot \frac{C}{2}$

#  $\tan\left(\frac{B-C}{2}\right) = \frac{b-c}{b+c} \cot \frac{A}{2}$

#  $\tan\left(\frac{C-A}{2}\right) = \frac{c-a}{c+a} \cot \frac{B}{2}$

Half Angle formula

#  $\sin \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{bc}}$

#  $\sin \frac{B}{2} = \sqrt{\frac{(s-c)(s-a)}{ac}}$

#  $\sin \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{ab}}$

#  $\cos \frac{A}{2} = \sqrt{\frac{s(s-a)}{bc}}$

#  $\cos \frac{B}{2} = \sqrt{\frac{s(s-b)}{ac}}$

#  $\cos \frac{C}{2} = \sqrt{\frac{s(s-c)}{ab}}$

#  $\tan \frac{A}{2} = \sqrt{\frac{(s-b)(s-c)}{s(s-a)}}$

#  $\tan \frac{B}{2} = \sqrt{\frac{(s-c)(s-a)}{s(s-b)}}$

#  $\tan \frac{C}{2} = \sqrt{\frac{(s-a)(s-b)}{s(s-c)}}$

Area of Triangle

#  $\Delta = \frac{1}{2} \cdot b \cdot h$


#  $\Delta = \frac{1}{2} ab \sin C = \frac{1}{2} bc \sin A = \frac{1}{2} ca \sin B$

#  $\Delta = \sqrt{s(s-a)(s-b)(s-c)}$

#  $\Delta = \frac{abc}{4R}$  (R is circumradius) #  $\Delta = r \times s$  (r is Inradius)

#  $\Delta = 2R^2 \sin A \sin B \sin C$

Inradius




#  $r = \frac{\Delta}{s}$

#  $r = (s-a) \tan \frac{A}{2}$   
#  $r = (s-b) \tan \frac{B}{2}$   
#  $r = (s-c) \tan \frac{C}{2}$


#  $r = 4R \sin \frac{A}{2} \sin \frac{B}{2} \sin \frac{C}{2}$

Circumradius



#  $R = \frac{abc}{4\Delta}$

Exradii




#  $r_1 = \frac{\Delta}{s-a} = s \tan \frac{A}{2} = 4R \sin \frac{A}{2} \cos \frac{B}{2} \cos \frac{C}{2}$

#  $r_2 = \frac{\Delta}{s-b} = s \tan \frac{B}{2} = 4R \cos \frac{A}{2} \sin \frac{B}{2} \cos \frac{C}{2}$

#  $r_3 = \frac{\Delta}{s-c} = s \tan \frac{C}{2} = 4R \cos \frac{A}{2} \cos \frac{B}{2} \sin \frac{C}{2}$


m-n cot Theorem




#  $(m+n) \cot \theta = m \cot \alpha - n \cot \beta$

#  $(m+n) \cot \theta = n \cot \beta - m \cot \alpha$


# Ellipse



$$PF_1 + PF_2 = QF_1 + QF_2$$



- AA' (major axis) = 2a
- BB' (minor axis) = 2b
- foci =  $(\pm ae, 0)$
- Directrix  $\rightarrow x = \pm \frac{a}{e}$
- $PF_1 + PF_2 = 2a$
- $b^2 = a^2(1 - e^2)$  /  $e = \sqrt{1 - \frac{b^2}{a^2}}$
- Vertices =  $(\pm a, 0)$
- Latus Rectum =  $\frac{2b^2}{a}$
- End of LR =  $(\pm ae, \pm \frac{b^2}{a})$



- [b > a]
- AA' (minor axis) = 2a
- BB' (major axis) = 2b
- foci  $\rightarrow (0, \pm be)$
- Directrix  $\rightarrow y = \pm \frac{b}{e}$
- $PF_1 + PF_2 = 2b$
- LR =  $\frac{2a^2}{b}$
- Ends of LR =  $(\pm \frac{a^2}{b}, \pm b)$

- Two ellipses are similar if they have equal eccentricity.
- Ellipse with axes || to coordinate axes and centre  $(h, k) \rightarrow \frac{(x-h)^2}{a^2} + \frac{(y-k)^2}{b^2} = 1$

Length of LR =  $(\text{minor axis})^2 / \text{major axis} = 2e(a - ae)$


Eq<sup>n</sup> of an Ellipse referred to two perpendicular lines.

$$L_1: a_1x + b_1y + c_1 = 0 \Rightarrow \frac{(a_1x + b_1y + c_1)^2}{\sqrt{a_1^2 + b_1^2}} + \frac{(b_2x - a_2y + c_2)^2}{\sqrt{a_2^2 + b_2^2}} = 1$$

- Centre at intersection point of  $L_1, L_2$
- Major axis is along  $L_2$  ( $a > b$ )

Position of a point w.r.t. an Ellipse  $(h, k)$   
 $\frac{x^2}{a^2} + \frac{y^2}{b^2} - 1 >, =, < 0 \rightarrow$  outside, on, inside.

Auxiliary Circle / Eccentric Angle



Aux circle:  $x^2 + y^2 = a^2$   
 $P(a \cos \theta, a \sin \theta)$   
 $\theta$  is called eccentric angle of point P

Properties of ellipse

- Area of ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is  $\pi ab$ .
- Ratio of area of any triangle PQR inscribed in ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  and that of triangle formed by corresponding points on the aux circle is  $\frac{1}{2}$ .
- Semi LR is HM of segments of focal chord.
- Circle describe on focal length as diameter always touches auxiliary circle.

Director circle



locus of poi of  $\perp$  tangents

Important Properties related to tangents

- Locus of feet of perpendiculars from foci upon any tangent is an auxiliary circle.
- Product of lengths of perpendiculars from foci upon any tangent of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$  is  $b^2$ .
- Tangents at the extremities of Latus Rectum pass through the corresponding foot of directrix on major axis.
- Length of tangent b/w the point of contact and the point where it meets the directrix subtends right angle at the corresponding focus.

Equation of tangent

- Point form  $\rightarrow \frac{xx_1}{a^2} + \frac{yy_1}{b^2} = 1$
- Parametric form  $\rightarrow \frac{x}{a} \cos \theta + \frac{y}{b} \sin \theta = 1$  ( $a \cos \theta, b \sin \theta$ )
- Slope form  $\rightarrow y = mx \pm \sqrt{a^2 m^2 + b^2}$

Eq<sup>n</sup> of Normal

- Point form  $\rightarrow \frac{a^2 x}{x_1} - \frac{b^2 y}{y_1} = a^2 - b^2$
- Parametric form  $\rightarrow ax \sec \theta - by \csc \theta = a^2 - b^2$

Properties of normals

- Normal other than major axis never passes through the focus.
- Normal at the point P bisects angle SPS' [Reflection property]

Co-normal Points: From any point in the plane maximum four normals can be drawn.


Eccentric angle of all the four points  $\alpha, \beta, \gamma, \delta$  then  $\alpha + \beta + \gamma + \delta = (2n+1)\pi$

Concyclic Points:  $\alpha + \beta + \gamma + \delta = 2n\pi$

Eq<sup>n</sup> of chord joining P( $\alpha$ ) & Q( $\beta$ )  $\rightarrow \frac{x}{a} \cos \frac{(\alpha+\beta)}{2} + \frac{y}{b} \sin \frac{(\alpha+\beta)}{2} = \cos \frac{(\alpha-\beta)}{2}$

POI of tangents at P( $\alpha$ ) & Q( $\beta$ )  $\rightarrow \left( a \frac{\cos \frac{\alpha+\beta}{2}}{\cos \frac{\alpha-\beta}{2}}, b \frac{\sin \frac{\alpha+\beta}{2}}{\cos \frac{\alpha-\beta}{2}} \right)$





$P_1F_1 - P_1F_2 = P_2F_1 - P_2F_2 = \text{const.}$   
 $LR = \frac{2b^2}{a} = 2c(ae - \frac{b^2}{a^2})$   
 Hyperbolas referred to two lines:

$L_1: lx + my + n = 0$   
 $L_2: mx - ly + p = 0$

$$\frac{(lx + my + n)^2}{a^2} - \frac{(mx - ly + p)^2}{b^2} = 1$$

- Centre is pol of  $L_1$  &  $L_2$
- TA  $\rightarrow 2a$ , CA  $\rightarrow 2b$
- TA is along  $L_2 = 0$

Equation of normal


- point form  $(x, y): \frac{ax}{x} + \frac{by}{y} = a^2 + b^2$
- parametric form:  $ax \cos \theta + by \cot \theta = a^2 + b^2$

Properties of Normals


- Normal other than TA never passes through focus.
- Locus of feet of perpendicular drawn from focus upon any tangent of hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is its aux circle i.e.  $x^2 + y^2 = a^2$
- The product of perpendiculars drawn from foci upon any tangent of the hyperbola  $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  is  $b^2$
- The portion of the tangent b/w the poc and the point where it meets the directrix subtends a right angle at corresponding focus.
- The tangent and normal at any point of ~~conjugate~~ hyperbola bisect the angle b/w focal radii.
- If an ellipse and a hyperbola have same foci, they cut at right angles.
- The foci and the points P & Q in which any tangent meets the tangents at the vertices are concyclic with PQ as diameter.

Hyperbola

Standard Eqn



Director Circle



- $a > b$ , DC is real
- $a = b$ , DC is point circle
- $a < b$ , no real circle

Chord with mp given  
 $T = S_1$


Conjugate Hyperbola

- Hyperbola  $\rightarrow \frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$  ( $e_1$ )
- conjugate Hyperbola  $\rightarrow \frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$  ( $e_2$ )
- $\frac{1}{e_1^2} + \frac{1}{e_2^2} = 1$
- foci of the hyperbola and conj. are concyclic square.

Eqn of Tangent


- Point form:  $\frac{xx_1}{a^2} - \frac{yy_1}{b^2} = 1$
- Parametric form:  $\frac{x}{a} \operatorname{sech} \theta - \frac{y}{b} \tanh \theta = 1$
- slope form:  $y = mx \pm \sqrt{a^2m^2 - b^2}$
- at  $(x, y)$  to  $\frac{(x-b)^2}{a^2} - \frac{(y-k)^2}{b^2} = 1$
- $\frac{(x-h)(x-k)}{a^2} - \frac{(y-l)(y-m)}{b^2} = 1$

Asymptotes



- Asymptotes pass thru the centre of the hyperbola
- The eqn of pair of asymptotes differ from the eqn of hyperbola just by only a constant.
- Asymptotes are diagonals of rectangle formed by lines drawn through extremities of ~~the~~ each axis parallel to the others.
- For rectangular hyperbola, Asymptotes are at  $90^\circ$  i.e.  $y = \pm x$
- At any point of a asymptote if a st. line is drawn perpendicular to the transverse axis, the product of the segments of this line, intercepted b/w the point and curve is always equal to the square of the semi conjugate axis.
- Perpendiculars from the foci on either asymptote meet it at the same point as the corresponding directrix and common points of intersection lie on aux circle.
- If the asymptotes of a rectangular hyperbola are  $x = \alpha$  and  $y = \beta$ , then its eqn is  $(x - \alpha)(y - \beta) = c^2$

Auxiliary Circle and Eccentric Angle



$Q(a \cos \theta, a \sin \theta)$   
 $P(a \operatorname{sech} \theta, b \tanh \theta)$

POJ of tangents from P(alpha) & Q(beta)

$$\left( a \frac{\cos(\frac{\alpha-\beta}{2})}{\cos(\frac{\alpha+\beta}{2})}, b \frac{\sin(\frac{\alpha+\beta}{2})}{\cos(\frac{\alpha-\beta}{2})} \right)$$

Eqn of chord joining P(alpha) and Q(beta)


$$\frac{x}{a} \cos(\frac{\alpha-\beta}{2}) - \frac{y}{b} \sin(\frac{\alpha+\beta}{2}) = \cos(\frac{\alpha+\beta}{2})$$

Pair of Tangents:  $SS_1 = T^2$

Important Points

- If angle b/w asymptotes of hyperbola is  $2\theta$ ,  $e = \sec \theta$
- Angle b/w asymptotes  $\theta = \tan^{-1} \left| \frac{2ab}{a^2 - b^2} \right|$
- Hyperbola and its conjugate have same asymptotes.

Rectangular Hyperbola



- Parametric form  $(ct, \frac{c}{t})$
- Equation of tangent at 't':  $x + yt^2 - 2ct = 0$
- Eqn of Normal at 't':  $xt^3 - yt - ct^4 + c = 0$
- Eqn of tangent at  $(x_1, y_1)$ :  $xy_1 + yx_1 = 2c^2$
- Eqn of normal at  $(x_1, y_1)$ :  $xx_1 - yy_1 = x_1^2 - y_1^2$

- $xy = c^2$ ,  $e = \sqrt{2}$
- Asymptotes,  $x=0, y=0$
- TA  $\neq y=x$ , CA:  $y=-x$
- Vertex:  $A(c, c)$ ,  $A'(-c, -c)$
- Foci  $(c\sqrt{2}, c\sqrt{2})$  &  $(-c\sqrt{2}, -c\sqrt{2})$
- length of LR  $= 2\sqrt{2}c$
- Aux circle  $\rightarrow x^2 + y^2 = c^2$
- DC  $\rightarrow x^2 + y^2 = 0$
- $x^2 - y^2 = 1$  and  $xy = 1$  intersect at  $90^\circ$

Concyclic points on  $xy = c^2$

If a circle and a rectangular hyperbola  $xy = c^2$  meet at four points  $t_1, t_2, t_3, t_4$ , then,

- $t_1 t_2 t_3 t_4 = 1$
- Centre of the mean position of the four points bisects the distance b/w the centres of the two curves.

# Theory of Equations and Logarithm

## ⊙ Laws of log

- $\log_a x = x \log_a a$ ;  $a, b > 0 \neq 1, x > 0$
- $\log_a x = \frac{1}{\log_x a}$
- $\log_a a = 1, \log_a 1 = 0$
- $\log_a x = \log_b x \cdot \log_a b = \frac{\log_b x}{\log_b a}$
- $\log_a (m^n) = n \log_a m$
- $\log_{a^n} (x) = \frac{1}{n} \log_a x$
- $\log_{a^n} x^m = \frac{m}{n} \log_a x$
- for  $x > y > 0$

(i)  $\log_a x > \log_a y$ , if  $a > 1$

(ii)  $\log_a x < \log_a y$ , if  $0 < a < 1$

•  $0 < a < 1$  then

(i)  $\log_a x > p \Rightarrow 0 < x < a^p$

(ii)  $\log_a x < p \Rightarrow a^p < x < 1$

•  $a > 1$ ,

(i)  $\log_a x > p \Rightarrow x > a^p$

(ii)  $0 < \log_a x < p \Rightarrow 0 < x < a^p$

## ⊙ Common Roots

• 1 common  $\rightarrow (D_{w_1})^2 = \text{Pass. Pass}$

• 2 common  $\rightarrow \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2}$

## ⊙ Relation b/w roots and Co-eff

$$a_0 x^n + a_1 x^{n-1} + a_2 x^{n-2} + \dots + a_{n-1} x + a_n = 0$$

•  $\sum \alpha_i = -\frac{a_1}{a_0}$  •  $\sum \alpha_1 \alpha_2 = \frac{a_2}{a_0}$  ..... •  $\alpha_1 \alpha_2 \alpha_3 \dots \alpha_n = (-1)^n \frac{a_n}{a_0}$

## ⊙ Discriminant & Nature of Roots

$$ax^2 + bx + c = 0 \rightarrow x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$






$$D = b^2 - 4ac$$

$D > 0 \rightarrow$  roots are real and distinct.

$D = 0 \rightarrow$  roots are real and equal

$D < 0 \rightarrow$  roots are imaginary.

•  $f(x) = y = ax^2 + bx + c$





## Binomial Theorem

- $(x+a)^n = {}^n C_0 x^n + {}^n C_1 x^{n-1} a + {}^n C_2 x^{n-2} a^2 + \dots + {}^n C_r x^{n-r} a^r + \dots + {}^n C_{n-1} x^1 a^{n-1} + {}^n C_n x^0 a^n$   
 → General Term:  $T_{r+1} = {}^n C_r x^{n-r} a^r$
- $(x-a)^n = {}^n C_0 x^n - {}^n C_1 x^{n-1} a + {}^n C_2 x^{n-2} a^2 - \dots + (-1)^r {}^n C_r x^{n-r} a^r + \dots + (-1)^n {}^n C_n x^0 a^n$   
 → General Term:  $T_{r+1} = (-1)^r {}^n C_r x^{n-r} a^r$
- Middle term: (i)  $(\frac{n}{2} + 1)$ th term, if  $n$  is even.  $T_{\frac{n}{2}+1} = {}^n C_{\frac{n}{2}} x^{\frac{n}{2}} a^{\frac{n}{2}}$   
 (ii)  $(\frac{n+1}{2})$ th &  $(\frac{n+3}{2})$ th term, if  $n$  is odd.

### Properties of Binomial Theorem

- ${}^n C_r = \frac{n}{r} {}^{n-1} C_{r-1} = \frac{n}{r} \cdot \frac{n-1}{r-1} \dots {}^{n-2} C_{r-2}$
- ${}^n C_r + {}^n C_{r-1} = {}^{n+1} C_r$
- ${}^n C_r = {}^n C_{n-r}$
- ${}^n C_0 + {}^n C_1 + {}^n C_2 + \dots + {}^n C_n = 2^n$
- ${}^n C_0 + {}^n C_2 + {}^n C_4 + \dots = {}^n C_1 + {}^n C_3 + \dots = 2^{n-1}$

• Greatest term  $\frac{T_{r+1}}{T_r} \geq 1$  i.e.  $\frac{n-r+1}{r} \left| \frac{a}{x} \right| \geq 1$

- $\sum_{r=0}^n (-1)^r {}^n C_r = 0$
- ${}^n C_1 - 2 {}^n C_2 + 3 {}^n C_3 - \dots + n (-1)^{n-1} {}^n C_n = 0$
- ${}^n C_1 + 2 {}^n C_2 + 3 {}^n C_3 + \dots + n {}^n C_n = n \cdot 2^{n-1}$
- ${}^n C_0 C_r + {}^n C_1 C_{r+1} + \dots + {}^n C_{n-r} C_n = 2^n C_{n-r}$
- ${}^n C_n + {}^{n+1} C_n + \dots + 2^{n-1} C_n = 2^n C_{n+1}$

### Multinomial Theorem

- $(x_1 + x_2 + \dots + x_k)^n = \sum_{r_1+r_2+\dots+r_k=n} \frac{n! x_1^{r_1} x_2^{r_2} \dots x_k^{r_k}}{r_1! r_2! \dots r_k!}$
- no of terms in  $(x+y+z)^n$  is  $n+2 C_2$  or  $\frac{(n+1)(n+2)}{2}$

### Expressions

- $(1+x)^{-1} = 1 - x + x^2 - x^3 + \dots + (-x)^r + \dots$
- $(1-x)^{-1} = 1 + x + x^2 + \dots + x^r + \dots$
- $(1+x)^{-2} = 1 - 2x + 3x^2 - 4x^3 + \dots + (r+1)(-x)^r + \dots$
- $(1-x)^{-2} = 1 + 2x + 3x^2 + 4x^3 + \dots + (r+1)x^r + \dots$
- $(1+x)^{-3} = 1 - 3x + 6x^2 - \dots + \frac{(r+1)(r+2)}{2!} (-x)^r + \dots$
- $(1-x)^{-3} = 1 + 3x + 6x^2 + \dots + \frac{(r+1)(r+2)}{2!} (x)^r + \dots$

# Determinants

$$\Delta = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \begin{matrix} \leftarrow R_1 \\ \leftarrow R_2 \\ \leftarrow R_3 \end{matrix}$$

[3x3]

## Minor

- Minor of  $a_{11}, M_{11} = \begin{vmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{vmatrix} = a_{22}a_{33} - a_{32}a_{23}$
- Minor of  $a_{21}, M_{21} = \begin{vmatrix} a_{11} & a_{13} \\ a_{31} & a_{33} \end{vmatrix} = a_{11}a_{33} - a_{31}a_{13}$
- Minor of  $a_{32}, M_{32} = \begin{vmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{vmatrix} = a_{11}a_{22} - a_{21}a_{12}$

## Co-factor

Co-factor of  $a_{ij} = (-1)^{i+j} M_{ij}$

Co-factor of  $a_{11} = (-1)^{1+1} M_{11} = M_{11}$

Co-factor of  $a_{12} = (-1)^{1+2} M_{12} = -M_{12}$

## Expansion of Determinants

$$\Delta = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} = a_{11}(a_{22}a_{33} - a_{32}a_{23}) - a_{12}(a_{21}a_{33} - a_{31}a_{23}) + a_{13}(a_{21}a_{32} - a_{22}a_{31})$$

## Properties

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = 0$$

any row/column

$$\begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} = - \begin{vmatrix} d & e & f \\ a & b & c \\ g & h & i \end{vmatrix}$$

Whenever we interchange any two row (or column) value of it will be multiplied by '-ve'

$$\begin{vmatrix} a+b & c & d \\ e+f & g & h \\ i+j & k & l \end{vmatrix} = \begin{vmatrix} a & c & d \\ e & g & h \\ i & k & l \end{vmatrix} + \begin{vmatrix} b & c & d \\ f & g & h \\ j & k & l \end{vmatrix}$$

$$\begin{vmatrix} ap & bp & cp \\ d & e & f \\ g & h & i \end{vmatrix} = p \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix}$$

Any two rows or column same.

Transform

$$\Delta = \begin{vmatrix} a & b & c \\ d & e & f \\ g & h & i \end{vmatrix} \xrightarrow{R_1 = R_1 + PR_2} \Delta = \begin{vmatrix} a+dp & b+ep & c+fp \\ d & e & f \\ g & h & i \end{vmatrix}$$

$$\begin{vmatrix} ap & d & g \\ bp & e & h \\ cp & f & i \end{vmatrix} = p \begin{vmatrix} a & d & g \\ b & e & h \\ c & f & i \end{vmatrix}$$

$$\Delta = \begin{vmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{vmatrix} \rightarrow a_{11}C_{11} + a_{12}C_{12} + a_{13}C_{13} = 0$$

or

$$a_{11}a_{31} + a_{12}C_{32} + a_{13}C_{33} = 0$$

or

$$a_{11}C_{13} + a_{22}C_{23} + a_{32}C_{33} = 0$$

## Important Expansion

$$\begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix} = (a-b)(b-c)(c-a) \cdot \begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^3 & b^3 & c^3 \end{vmatrix} = (a-b)(b-c)(c-a)(a+b+c) \cdot \begin{vmatrix} 1 & 1 & 1 \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{vmatrix} = (a-b)(b-c)(c-a)(ab+bc+ca)$$

$$\begin{vmatrix} a & b & c \\ b & c & a \\ c & a & b \end{vmatrix} = (a+b+c)(a^2+b^2+c^2-ab-bc-ca)$$

## System of Evaluate

$$\begin{cases} a_1x + b_1y + c_1z = d_1 \\ a_2x + b_2y + c_2z = d_2 \\ a_3x + b_3y + c_3z = d_3 \end{cases}$$

### Note

- If  $[\Delta \neq 0]$ 
  - Unique Sol<sup>n</sup>
  - Consistent set of sol<sup>n</sup>

$$\Delta = \begin{vmatrix} a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \\ a_3 & b_3 & c_3 \end{vmatrix}$$

- If  $[\Delta = 0]$

→ (a) if any one (or two)

$$\Delta_x = \begin{vmatrix} d_1 & b_1 & c_1 \\ d_2 & b_2 & c_2 \\ d_3 & b_3 & c_3 \end{vmatrix}$$

of  $\Delta_x, \Delta_y, \Delta_z$  is/one Non-zero, Inconsistent System  $\Rightarrow$  No sol<sup>n</sup>

$$\Delta_y = \begin{vmatrix} d_1 & a_1 & c_1 \\ d_2 & a_2 & c_2 \\ d_3 & a_3 & c_3 \end{vmatrix}$$

→ (b)  $[\Delta_x = \Delta_y = \Delta_z = 0]$

$$\Delta_z = \begin{vmatrix} d_1 & b_1 & a_1 \\ d_2 & b_2 & a_2 \\ d_3 & b_3 & a_3 \end{vmatrix}$$

Consistent set of sol<sup>n</sup>  $\Rightarrow$  Infinite no of sol<sup>n</sup>

$$\begin{cases} x = \frac{\Delta_x}{\Delta} \\ y = \frac{\Delta_y}{\Delta} \\ z = \frac{\Delta_z}{\Delta} \end{cases} \text{Cramer's Rule } (\Delta \neq 0)$$



## Trigonometric Equation

•  $\sin \theta = 0 \longrightarrow \theta = n\pi, n \in \mathbb{Z}$

•  $\cos \theta = 0 \longrightarrow \theta = (2n+1)\frac{\pi}{2}, n \in \mathbb{Z}$

•  $\tan \theta = 0 \longrightarrow \theta = n\pi, n \in \mathbb{Z}$

•  $\sin \theta = 1 \longrightarrow \theta = (4n+1)\frac{\pi}{2}, n \in \mathbb{Z}$

•  $\sin \theta = -1 \longrightarrow \theta = (4n-1)\frac{\pi}{2}, n \in \mathbb{Z}$

•  $\cos \theta = 1 \longrightarrow \theta = 2n\pi, n \in \mathbb{Z}$

•  $\cos \theta = -1 \longrightarrow \theta = (2n+1)\pi, n \in \mathbb{Z}$

•  $\cot \theta = 0 \longrightarrow \theta = (2n+1)\frac{\pi}{2}, n \in \mathbb{Z}$

•  $\sin \theta = \sin \alpha \longrightarrow n\pi + (-1)^n \alpha, n \in \mathbb{Z}$

$\Rightarrow \sin \theta = k \longrightarrow \theta = n\pi + (-1)^n (\sin^{-1} k), n \in \mathbb{Z}$

•  $\cos \theta = \cos \alpha \longrightarrow \theta = 2n\pi \pm \alpha, n \in \mathbb{Z}$

$\Rightarrow \cos \theta = k \longrightarrow \theta = 2n\pi \pm (\cos^{-1} k), n \in \mathbb{Z}, k \in [-1, 1]$

•  $\tan \theta = \tan \alpha \longrightarrow \theta = n\pi + \alpha, n \in \mathbb{Z}$

$\Rightarrow \tan \theta = k \longrightarrow \theta = n\pi + (\tan^{-1} k), k \in \mathbb{R}$

•  $\sin^2 \theta = \sin^2 \alpha / \cos^2 \theta = \cos^2 \alpha$

$\longrightarrow \theta = n\pi \pm \alpha, n \in \mathbb{Z}$

•  $\tan^2 \theta = \tan^2 \alpha \longrightarrow \theta = n\pi \pm \alpha, n \in \mathbb{Z}$

• Solution of the equation of the form  $a \cos \theta + b \sin \theta = c$

$\rightarrow$  If  $|c| > \sqrt{a^2 + b^2}$ , then no real solution

$\rightarrow$  If  $|c| \leq \sqrt{a^2 + b^2}$ , then divide both sides of the equation

by  $\sqrt{a^2 + b^2}$ , then take  $\cos \alpha = \frac{a}{\sqrt{a^2 + b^2}}$ ,

$\sin \alpha = \frac{b}{\sqrt{a^2 + b^2}}$ , equation will reduce to

$\cos(\theta - \alpha) = \cos \beta$ , where  $\tan \alpha = \frac{b}{a}$

$\cos \beta = \frac{c}{\sqrt{a^2 + b^2}}$

\* If we take  $\sin \alpha = \frac{a}{\sqrt{a^2 + b^2}}$ ,

$\cos \alpha = \frac{b}{\sqrt{a^2 + b^2}}$ , then the equation will

reduce to  $\sin(\theta + \alpha) = \sin \beta$ ,

$\sin \beta = \frac{c}{\sqrt{a^2 + b^2}}$

$\Rightarrow$  While solving triago equation, avoid squaring the equation as far as possible. If squaring is necessary check the solution for extraneous values (similar values following the same pattern).

$\Rightarrow$  Never cancel terms containing unknown terms on the two sides which are in product. It may cause the loss of a genuine solution.

$\Rightarrow$  The answer should not contain such values of angles which make any term undefined or infinite.



$\Rightarrow$  Domain should not change while simplifying the equation. If it changes, necessary corrections must be made.

$\Rightarrow$  check the denominator is not zero at any stage while solving the equation.


• Extreme values of functions ↖ keep in mind.

# Functions





Condition: A mapping is a function in each ~~each~~ input have one and only one outputs.



If no vertical line cuts the graph more than once, it's a function




Types of mapping/function:





- one-one  $\rightarrow$  Injection
- Onto  $\rightarrow$  Surjection
- one-one-onto  $\rightarrow$  Bijection

Slope:  
 strictly increasing ( $\frac{dy}{dx} > 0$ )  
 strictly decreasing ( $\frac{dy}{dx} < 0$ )
















If a horizontal line cuts the graph at more than one point, it's a many-one or else one-one

Type of function:



Fundamental graphs:



Signum function:  
 $\text{sgn}(x) = \frac{|x|}{x} = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$

Graphical Transformation:  
 $y = f(x) \rightarrow y = -f(x)$ : mirror img about x axis  
 $y = f(x) \rightarrow y = f(x) \pm k$ :  $+k \rightarrow$  up /  $-k \rightarrow$  down  
 $y = f(x) \rightarrow y = f(-x)$ : mirror img about y axis  
 $y = f(x) \rightarrow y = |f(x)|$   
 $y = f(x) \rightarrow y = f(|x|)$

min/max function:  
 Graph showing a curve with a local maximum.

Inverse function:  
 $f(x)$  is invertible only if it is one-one-onto / Bijective.  
 $f^{-1}(f(x)) = x$  |  $[f'(f(x))] f'(x) = 1$  |  $f^{-1}(f(x))' = \frac{1}{f'(x)}$  Put  $x=1, y=k$  (as)

Periodic function: If  $f(x+T) = f(x), T > 0$   $f(x)$  is called period function with period  $T$ .  
 smallest value of  $T$  is called fundamental period of  $f(x)$

$\rightarrow$  period of  $\sin x, \cos x, \csc x, \sec x \rightarrow 2\pi$  |  $\tan x, \cot x \rightarrow \pi$  |  $|\sin x|, |\cos x| \dots \rightarrow \pi$   
 $\rightarrow \sin^n x, \cos^n x, \csc^n x, \sec^n x \rightarrow \begin{cases} 2\pi, & n \text{ odd} \\ \pi, & n \text{ even} \end{cases}$  |  $\tan^n x, \cot^n x \rightarrow \pi$  | if  $f(x) \rightarrow T \rightarrow f(x) \pm k \rightarrow T$   
 $\rightarrow$  const functions are periodic but period not defined. |  $f(x) \rightarrow T_1, g(x) \rightarrow T_2 \rightarrow f(x) \pm g(x) / \frac{f(x)}{g(x)} \rightarrow \text{LCM}(T_1, T_2)$   
 $f(x) \cdot g(x) \rightarrow \text{LCM}(T_1, T_2)$

$\rightarrow f(x-a) = f(x+a) \rightarrow T = 2a$   
 $\rightarrow f(a-x) = f(a+x) \rightarrow f(x)$  is symmetric abt. period  $2(a-x)$   
 $\rightarrow f(a-x) = f(a+x)$  &  $f(b-x) = f(b+x) \rightarrow$  periodic abt.  $2(a-b)$   
 $\rightarrow f(a-x) = f(a+x) \rightarrow 2|ab-a| \rightarrow 2|a-c|$   
 $f(b-x) = f(b+x) \rightarrow 2|b-c|$   
 $f(c-x) = f(c+x) \rightarrow 2|c-a|$   
 Period of  $f(x) = \min \{ 2|a-b|, 2|b-c|, 2|c-a| \}$



# Limits


## Expansions:

- $\sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \frac{x^7}{7!} + \dots$
- $\cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \frac{x^6}{6!} + \dots$
- $\tan x = x + \frac{x^3}{3} + \frac{2}{15}x^5 + \dots$
- $\log(1+x) = x - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$
- $\log(1-x) = -x - \frac{x^2}{2} - \frac{x^3}{3} - \frac{x^4}{4} - \dots$
- $\log e^x = 1 + \frac{x}{1!} + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots$
- $a^x = 1 + \frac{x \ln a}{1!} + \frac{(x \ln a)^2}{2!} + \frac{(x \ln a)^3}{3!} + \dots$
- $\tan^{-1} x = x - \frac{x^3}{3} + \frac{x^5}{5} - \frac{x^7}{7} + \dots$
- $\sin^{-1} x = x + \frac{x^3}{6} + \frac{3}{40}x^5 + \dots$


## Important Results

- $\lim_{x \rightarrow 0} \frac{\sin x}{x} = \lim_{x \rightarrow 0} \frac{x}{\sin x} = \lim_{x \rightarrow 0} \frac{\tan x}{x} = \lim_{x \rightarrow 0} \frac{x}{\tan x} = 1$
- $\lim_{x \rightarrow 0} \frac{\sin^{-1} x}{x} = \lim_{x \rightarrow 0} \frac{\tan^{-1} x}{x} = 1$
- $\lim_{x \rightarrow 0} \frac{e^x - 1}{x} = 1$
- $\lim_{x \rightarrow a} \frac{a^x - 1}{x} = \ln a$
- $\lim_{x \rightarrow a} \frac{x^n - a^n}{x - a} = na^{n-1}$
- $\lim_{x \rightarrow 0} \frac{\ln(1+x)}{x} = 1$


## Continuity



If  $\lim_{x \rightarrow a^-} f(x) = \lim_{x \rightarrow a^+} f(x) = f(a)$   
 $\rightarrow y = f(x)$  is continuous at  $x = a$




If  $\lim_{x \rightarrow a} f(x) \neq f(a)$ , Discontinuous at  $x = a$ , point discontinuity / Removable discontinuity.




$LHL \neq RHL$ , Discontinuous at  $x = a$ .  
 Jump Discontinuity.

## Differentiability



If  $LHD = RHD$  at  $x = a$ ,  $f(x)$  is differentiable at  $x = a$

- Sharp turns lead to non-differentiable points.
- Smooth curves are generally differentiable at all points.
- Tangents must have finite slope to make function differentiable.



RHD at  $x = a$

$$Rf'(a) = \lim_{h \rightarrow 0} \frac{f(a+h) - f(a)}{h}$$

LHD at  $x = a$


$$Lf'(a) = \lim_{h \rightarrow 0} \frac{f(a-h) - f(a)}{-h}$$

Discontinuous  $\Rightarrow$  non-differentiable

Differentiable  $\Rightarrow$  continuous.

$f(x) \rightarrow \text{diff} \rightarrow f'(x) \rightarrow \text{cont} / f''(x) \rightarrow \text{cont} \rightarrow f'(x) \rightarrow \text{diff}$

# Application of Derivatives



- Slope of tangent =  $\frac{d}{dx} f(x) |_{(x_1, y_1)} = \tan \theta$
- Slope of normal =  $-\frac{dx}{dy} |_{(x_1, y_1)} = -\cot \theta$
- Equation of tangent:  $(y - y_1) = \frac{dy}{dx} |_{(x_1, y_1)} (x - x_1)$
- Equation of normal:  $(y - y_1) = -\frac{dx}{dy} |_{(x_1, y_1)} (x - x_1)$

length of tangent (PT) =  $|y_1| \operatorname{cosec} \theta$  ,  $|PT| = |y_1| \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$


length of normal (PN) =  $|y_1| \operatorname{sec} \theta$  ,  $|PN| = |y_1| \sqrt{1 + \left(\frac{dy}{dx}\right)^2}$

length of Sub-tangent =  $|y_1| \cot \theta$  ,  $|TM| = |y_1| \left| \frac{dx}{dy} \right|$

length of Sub-normal =  $|y_1| \tan \theta$  ,  $|MN| = |y_1| \left| \frac{dy}{dx} \right|$

## Rolle's Theorem

$f(x)$  is cont on  $[a, b]$ , diff on  $(a, b)$  and  $f(a) = f(b)$   
 Then there exist at least one  $c \in (a, b)$  so that  $f'(c) = 0$



## Maxima/Minima:

$\frac{dy}{dx} = 0 \Rightarrow x = a, b$



$\frac{d^2y}{dx^2} = \frac{d}{dx} \left( \frac{dy}{dx} \right)$

$\frac{d^2y}{dx^2} |_{x=a} < 0$  [ $x=a$  is maxima]


$\frac{d^2y}{dx^2} |_{x=b} > 0$  [ $x=b$  is minima]

$\frac{d^2y}{dx^2} |_{x=c} = 0$  [ $x=c$  is inflection]

## Critical points:



## Angle b/w 2 curves



$\frac{d}{dx} f(x) |_{(x_1, y_1)} = m_1$

$\frac{d}{dx} g(x) |_{(x_1, y_1)} = m_2$


$\theta = 90^\circ$  is called orthogonal intersection.

$\theta = \tan^{-1} \left| \frac{m_1 - m_2}{1 + m_1 m_2} \right|$

## Lagrange's Mean Value Theorem (LMVT)


if  $y = f(x)$  is continuous on  $[a, b]$  & differentiable on  $(a, b)$ , There exist at least one such value of  $c \in (a, b)$  such that

$f'(c) = \frac{f(b) - f(a)}{b - a}$




- Monotonicity:  $\frac{dy}{dx} > 0 \Rightarrow y = f(x)$  is an increasing function
- $\frac{dy}{dx} < 0 \Rightarrow y = f(x)$  is a strictly decreasing function
- $\frac{dy}{dx} \geq 0 \Rightarrow y = f(x)$  is a non-decreasing function
- $\frac{dy}{dx} \leq 0 \Rightarrow y = f(x)$  is a non-increasing function.

Monotonicity/Maxima/Minima have NOTHING to do with continuity of the graph.



$f'(x) = (x-1)^2(x-2)x^2(x+1)^5$




Note:  $\frac{d^2y}{dx^2} = 0$  at  $x=a$  is a point of inflection provided  $\frac{d^3y}{dx^3}$  is non zero at  $x=a$



Indefinite integrals


- $\frac{d}{dx} x^n = nx^{n-1} \rightarrow \int x^n dx = \frac{x^{n+1}}{n+1} + c$
- $\frac{d}{dx} \log x = \frac{1}{x} \rightarrow \int \frac{1}{x} dx = \log x + c$
- $\frac{d}{dx} e^x = e^x \rightarrow \int e^x dx = e^x$
- $\frac{d}{dx} a^x = a^x \ln a \rightarrow \int a^x dx = \frac{a^x}{\ln a} + c$
- $\frac{d}{dx} \sin x = \cos x \rightarrow \int \cos x dx = \sin x + c$
- $\frac{d}{dx} \cos x = -\sin x \rightarrow \int \sin x dx = -\cos x + c$
- $\frac{d}{dx} \tan x = \sec^2 x \rightarrow \int \sec^2 x dx = \tan x + c$
- $\frac{d}{dx} \cot x = -\csc^2 x \rightarrow \int \csc^2 x dx = -\cot x + c$
- $\frac{d}{dx} \sec x = \sec x \tan x \rightarrow \int \sec x \tan x dx = \sec x + c$
- $\frac{d}{dx} \csc x = -\csc x \cot x \rightarrow \int \csc x \cot x dx = -\csc x + c$
- $\frac{d}{dx} \sin^{-1} x = \frac{1}{\sqrt{1-x^2}} \rightarrow \int \frac{1}{\sqrt{1-x^2}} dx = \sin^{-1} x + c$
- $\frac{d}{dx} \tan^{-1} x = \frac{1}{1+x^2} \rightarrow \int \frac{1}{1+x^2} dx = \tan^{-1} x + c$
- $\frac{d}{dx} \sec^{-1} x = \frac{1}{x\sqrt{x^2-1}} \rightarrow \int \frac{1}{x\sqrt{x^2-1}} dx = \sec^{-1} x + c$

- $\int \frac{1}{\sqrt{a^2-x^2}} dx = \sin^{-1}(\frac{x}{a}) + c$
- $\int \frac{1}{a^2+x^2} dx = \frac{1}{a} \tan^{-1}(\frac{x}{a}) + c$
- $\int \frac{1}{x\sqrt{x^2-a^2}} dx = \frac{1}{a} \sec^{-1}(\frac{x}{a}) + c$
- $\int \frac{1}{x-a} dx = \frac{1}{a} \log|\frac{x-a}{x+a}| + c$
- $\int \tan x dx = \log|\sec x| + c$
- $\int \cot x dx = \log|\sin x| + c$
- $\int \sec x dx = \log|\sec x + \tan x| + c$
- $\int \csc x dx = \log|\tan(\frac{\pi}{4} + \frac{x}{2})| + c$
- $\int \csc x \cot x dx = \log|\csc x - \cot x| + c$
- $\int \csc x dx = \log|\tan \frac{x}{2}| + c$




- By Parts:**  $\int I \cdot II dx = I \int II dx - \int (\frac{d}{dx} I) (\int II dx) dx$
- $\int e^x (f(x) + f'(x)) dx = e^x f(x) + c$
- Forms**
  - $\int \frac{1}{\text{linear}} dx = \frac{\log|\text{linear}|}{\text{coeff of } x} + c$
  - $\int \frac{1}{(\text{linear})^n} dx = \int \frac{(\text{linear})^{n+1}}{(-n+1)(\text{coeff of } x)} + c$
  - $\int \log x dx = x \log x - x + c$
  - $\int \frac{f'(x)}{f(x)} dx = \ln|f(x)| + c$
  - $\int (f(x))^n f'(x) dx = \frac{(f(x))^{n+1}}{n+1} + c$
  - $\int \frac{dx}{\text{Quad}}$
  - $\int \frac{dx}{a^2+x^2} = \frac{1}{a} \tan^{-1}(\frac{x}{a})$
  - $\int \frac{dx}{x^2-a^2} = \frac{1}{2a} \ln|\frac{x-a}{x+a}| + c$
  - $\int \frac{1}{a^2-x^2} dx = \frac{1}{2a} \ln|\frac{a+x}{a-x}| + c$
- $\int \frac{1}{a \sin x + b \cos x} dx \rightarrow$  put  $\sin x = \frac{2 \tan \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}$ ,  $\cos x = \frac{1 - \tan^2 \frac{x}{2}}{1 + \tan^2 \frac{x}{2}}$
- $\int \sin^m x \cos^n x dx$  ( $m, n \in \mathbb{N}$ )
  - If  $m, n \in \text{odd}$ , subs any
  - If one is odd, sub even
  - If both are even, use trigo
  - If both are rational and  $\frac{m+n-2}{2}$  is -ve int. then sub  $\cot x = p$  or  $\tan x = p$
- $\int \frac{dx}{a \cos^2 x + b \sin^2 x}$ ,  $\int \frac{dx}{a \sin^2 x}$ ,  $\int \frac{dx}{a \cos^2 x + b \sin^2 x}$ 
  - divide  $N^r$  &  $D^r$  by  $\cos^2 x$
- $\int \frac{p \cos x + q \sin x + r}{a \cos x + b \sin x + c} dx \rightarrow wR, N^r = \lambda(D^r) + \mu(\frac{dD^r}{dx}) + \gamma$
- Biquadratic**  $\rightarrow$  sub  $(x + \frac{1}{x})$  or  $(x - \frac{1}{x}) = t$
- $\int \frac{px+q}{ax^2+bx+c} dx$ ,  $\int \frac{px+a}{\sqrt{ax^2+bx+c}} dx \rightarrow wR px+q = \frac{d}{dx}(ax^2+bx+c) + \mu$
- $\int \frac{1}{L\sqrt{L_2}} dx$ ,  $\int \frac{L_1}{\sqrt{L_2}} dx$ ,  $\int \frac{\sqrt{L_2}}{L_1} dx \rightarrow$  sub  $L_2 = t^2$
- $\int \frac{1}{L\sqrt{L_2}} dx \rightarrow$  sub  $\frac{1}{L} = t$
- $\int \frac{1}{a_1\sqrt{a_2}} dx \rightarrow x = \frac{1}{t} \rightarrow$  Integrand will become  $\int \frac{t dt}{(p^2+t^2)(r^2+t^2)}$  then  $u^2 = r^2+t^2$
- $\int \sqrt{\text{Quad}} dx \rightarrow$ 
  - $\int \sqrt{a^2+x^2} dx = \frac{x}{2} \sqrt{a^2+x^2} + \frac{a^2}{2} \ln|x + \sqrt{a^2+x^2}| + c$
  - $\int \sqrt{x^2-a^2} dx = \frac{x}{2} \sqrt{x^2-a^2} - \frac{a^2}{2} \ln|x + \sqrt{x^2-a^2}| + c$
  - $\int \sqrt{a^2-x^2} dx = \frac{x}{2} \sqrt{a^2-x^2} + \frac{a^2}{2} \sin^{-1}(\frac{x}{a}) + c$

# DEFINITE INTEGRATIONS



$$\int_a^b f(x) dx = [F(x)]_a^b = F(b) - F(a)$$

Region lying above x axis will give +ve value of integral & negative for the portion lying below x axis.



$$\int_a^b f(x) dx = A_1 - A_2 + A_3$$

Properties:

$$\int_a^b f(x) dx = \int_a^b f(t) dt$$

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx$$

[c may or may not belong to (a,b)]

$$\int_a^b f(x) dx = \int_a^b f(a+b-x) dx$$

[Turning Property]

$$\int_{-a}^a f(x) dx = \begin{cases} 2 \int_0^a f(x) dx, & \text{if } f(x) \text{ is even func i.e. } f(-x) = f(x) \\ 0, & \text{if } f(x) \text{ is odd func} \end{cases}$$

While substituting, t must be continuous in the interval

Properties related to periodic func: [if  $f(x+T) = f(x)$ , period is T]

$$\int_0^{nT} f(x) dx = n \int_0^T f(x) dx, n \in \mathbb{I}$$

$$\int_m^{nT} f(x) dx = (n-m) \int_0^T f(x) dx, n, m \in \mathbb{I}$$

$$\int_{a+nT}^{a+nT+T} f(x) dx = \int_a^{a+T} f(x) dx$$





$$\int_0^{\pi/2} \log \sin x dx = \int_0^{\pi/2} \log \cos x dx = -\frac{\pi}{2} \log 2$$

Newton-Leibnitz Rule:

$$\frac{d}{dx} \left( \int_{h(x)}^{g(x)} h(t) dt \right) = h(g(x)) \times \frac{d}{dx}(g(x)) - h(h(x)) \times \frac{d}{dx}(h(x))$$

Leibnitz 2nd Rule:

$$\text{If } I(d) = \int_a^b f(x, d) dx \rightarrow \frac{\partial I}{\partial d} = \int_a^b \frac{\partial f(x, d)}{\partial d} dx$$



Vertical Strip:

$$\text{Area} = \int_{x=a}^{x=b} (\text{upper } y - \text{lower } y) dx$$

Horizontal Strip:

$$\text{Area} = \int_{y=c}^{y=d} (\text{Right } x - \text{left } x) dy$$



# DIFFERENTIAL EQUATION

- Eq involving  $x, y$  & differentials co-efficient. DE represents a family of curves.
- Order:** Order of highest order derivative present in the eq<sup>n</sup> is the order of D.E.
- Degree:** Degree of the highest order derivative present in the eq<sup>n</sup> is the degree of DE, provided the eq<sup>n</sup> is polynomial in different co-eff and eq<sup>n</sup> is free from radicals.

**Formation of DE:** (Degree of a DE = No of arbitrary constants present in eq<sup>n</sup>)

DE of all lines passing thru origin:  $y = mx$   $y = \frac{dy}{dx} x \rightarrow x dy - y dx = 0$  DE of all lines:  $y = mx + c$   
 $\frac{dy}{dx} = m$   $\frac{dy}{dx} = m$  ,  $\frac{d^2y}{dx^2} = 0$

**Solution of DE:**

**Variable-seperable form:**  $\frac{dy}{dx} = e^{x-y} + x^2 e^{-y}$  Eq<sup>n</sup> Reducible to Variable Seperable form

$\Rightarrow \frac{dy}{dx} = \frac{e^x + x^2}{e^y} \Rightarrow \int e^y dy = \int (e^x + x^2) dx$

$\frac{dy}{dx} = f(ax + by + c)$ , consider,  $ax + by + c = t$

**Homogeneous form:**

$\frac{dy}{dx} = \frac{f(x, y)}{g(x, y)}$  where  $f$  and  $g$  are of same order.

**Eq<sup>n</sup> reducible to Homogenous form:**  $\frac{dy}{dx} = \frac{ax + by + c}{Ax + By + D}$

[If  $aB \neq Ab$  or  $A + b \neq 0$ ]

$x = X + h$   $y = Y + k$

$dx = dX$   $dy = dY$

$\therefore \frac{dY}{dX} = \frac{aX + bY + ah + bk + c}{AX + BY + Ah + Bk + D}$

$ah + bk + c = 0$   
&  
 $Ah + Bk + D = 0$  } find value of  $h$  &  $k$

**Linear Differential Eq<sup>n</sup>:**

$\Rightarrow \frac{dy}{dx} + Py = Q$  [P & Q are func of  $x$  alone]

$\frac{dY}{dX} = \frac{aX + bY}{AX + BY}$   $\rightarrow$  Homogeneous

I.F. =  $e^{\int P dx}$

$\rightarrow y(I.F.) = \int Q(I.F.) dx$

• If  $aB = Ab$   $\rightarrow$  assume  $(ax + by = t)$

• If  $A + b = 0$   $\rightarrow$  simply cross multiply & replace  $x dy + y dx$  by  $d(xy)$

In the end,  $X = x - h$   
 $Y = y - k$

$\Rightarrow \frac{dx}{dy} + Mx = N$  [M & N are func of  $y$  alone]

**Bernoulli Eq<sup>n</sup>**

$\frac{dy}{dx} + \frac{y}{x} = y^n$

divide by  $y^n$  and then assume  $\frac{1}{y^{n-1}}$ , co-eff of  $x$  as  $t$ .

I.F. =  $e^{\int M dy}$

$\rightarrow x(I.F.) = \int (N(I.F.)) dy$

here,  $t = \frac{1}{y^{n-1}}$

# VECTORS

## Angle bisector b/w two vectors:

Internal  $\rightarrow \vec{R} = \lambda(\hat{a} + \hat{b})$


External  $\rightarrow \vec{Q} = \mu(\hat{a} - \hat{b})$

## Section formula:

Internal  $\rightarrow \left(\frac{m\vec{b} + n\vec{a}}{m+n}\right)$

External  $\rightarrow \left(\frac{m\vec{b} - n\vec{a}}{m-n}\right)$

## Dot (Scalar) Product:



$\vec{a} \cdot \vec{b} = |\vec{a}| |\vec{b}| \cos \theta$

$\vec{a} \cdot \vec{b} = a_1 b_1 + a_2 b_2 + a_3 b_3$

$\vec{a} \cdot \vec{a} = |\vec{a}|^2$

$\vec{a} \cdot \vec{b} = 0 \rightarrow$  perpendicular

$\hat{i} \cdot \hat{i} = \hat{j} \cdot \hat{j} = \hat{k} \cdot \hat{k} = 1$

$\hat{i} \cdot \hat{j} = \hat{j} \cdot \hat{k} = \hat{k} \cdot \hat{i} = 0$

## Angle b/w the vectors $\rightarrow$

$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|}$

Projection of  $\vec{a}$  on  $\vec{b} \rightarrow$   
 $\vec{p} = \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|}$


## Vector Triple Product:

$(\vec{a} \times \vec{b}) \times \vec{c} = (\vec{a} \cdot \vec{c}) \vec{b} - (\vec{b} \cdot \vec{c}) \vec{a}$

$\vec{a} \times (\vec{b} \times \vec{c}) = (\vec{a} \cdot \vec{c}) \vec{b} - (\vec{a} \cdot \vec{b}) \vec{c}$

## Cross (Vector) Product:

$\vec{a} \times \vec{b} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \end{vmatrix}$



$\text{Area} = \frac{1}{2} |\vec{a} \times \vec{b}|$

$\vec{a} \times \vec{b} = |\vec{a}| |\vec{b}| \sin \theta \hat{r}$

$\text{Area} = |\vec{a} \times \vec{b}|$

$\text{Area} = \frac{1}{2} |\vec{a} \times \vec{b} + \vec{b} \times \vec{c} + \vec{c} \times \vec{a}|$



## Scalar Triple product (Box Product):


$[\vec{a} \ \vec{b} \ \vec{c}] = \vec{a} \cdot (\vec{b} \times \vec{c}) \rightarrow$  volume of parallelepiped

$[k\vec{a} \ \vec{b} \ \vec{c}] = k[\vec{a} \ \vec{b} \ \vec{c}]$

$[\vec{a} \ \vec{b} \ \vec{c}] = [\vec{b} \ \vec{c} \ \vec{a}] = [\vec{c} \ \vec{a} \ \vec{b}]$

$[\vec{a} \ \vec{b} \ \vec{c}] = -[\vec{b} \ \vec{a} \ \vec{c}]$

$[\vec{a} \ \vec{b} \ \vec{c}] = 0$  if  $\vec{a}, \vec{b}, \vec{c}$  are coplanar.



volume of parallelepiped

$[\vec{a} \ \vec{b} \ \vec{c}] = \begin{vmatrix} a_1 & a_2 & a_3 \\ b_1 & b_2 & b_3 \\ c_1 & c_2 & c_3 \end{vmatrix}$

of all lines:  $y = mx + c$   
 degree of DE, provided the

$\frac{dy}{dx} = m, \frac{d^2y}{dx^2} = 0$

Separable form:  
 ax + by + c = t

$ax + by + c = 0$   
 $Ax + By + D = 0$

find value of h & k  
 $x + D = 0$


In the end,  $X = x - h$   
 $Y = y - k$

replace  $x dy + y dx$  by  $d(xy)$

coeff of x as t



**Direction Cosines:**



$l^2 + m^2 + n^2 = 1$


**Direction Ratios:** Simple ratio of DC.

DR  $\rightarrow$  DC  $\rightarrow$  DR  $(a, b, c) \rightarrow$  DC  $(\frac{a}{\sqrt{a^2+b^2+c^2}}, \frac{b}{\sqrt{a^2+b^2+c^2}}, \frac{c}{\sqrt{a^2+b^2+c^2}})$   
 DC  $\rightarrow$  DR  $\rightarrow$  DC  $(\frac{1}{\sqrt{3}}, -\frac{1}{\sqrt{3}}, \frac{1}{\sqrt{3}}) \rightarrow$  DR  $(1, -1, 1)$  or  $(2, -2, 2)$  or  $(\lambda, -\lambda, \lambda)$   
 $\rho(a_1, b_1, c_1) \& \rho(a_2, b_2, c_2) \rightarrow \cos \theta = \frac{a_1 a_2 + b_1 b_2 + c_1 c_2}{\sqrt{a_1^2 + b_1^2 + c_1^2} \sqrt{a_2^2 + b_2^2 + c_2^2}} = 0$  if  $\theta = 90^\circ$   
 $(l_1, m_1, n_1) \& (l_2, m_2, n_2) \rightarrow \cos \theta = l_1 l_2 + m_1 m_2 + n_1 n_2$

- $(l_1, m_1, n_1) \& (l_2, m_2, n_2)$  DC of 2 vectors  $\rightarrow$  internal bisector  $(l_1 + l_2, m_1 + m_2, n_1 + n_2)$ , external bisector  $(l_1 - l_2, m_2 - m_1, n_2 - n_1)$
- DR of line joining  $A(a_1, b_1, c_1)$  &  $B(a_2, b_2, c_2) \rightarrow (a_1 - a_2, b_1 - b_2, c_1 - c_2)$
- $x=0$  yz plane,  $y=0$  xz plane,  $z=0$  xy plane,  $x=y=z=0$  origin
- In 3D line is the intersection of 2 planes.

**Equation of line in 3D:**


Equation of line passing thru a point  $\vec{a}$  and parallel to another vector  $\vec{b}$



$\vec{r} = \vec{a} + \lambda \vec{b}$  vector form

$\vec{r} = (x, y, z), \vec{a} = (x_1, y_1, z_1), \vec{b} = (a, b, c)$   
 $\vec{r} - \vec{a} = \lambda \vec{b}$   
 $\frac{x-x_1}{a} = \frac{y-y_1}{b} = \frac{z-z_1}{c} = \lambda$  Cartesian form

Equation of line passing thru 2 points.



$\vec{r} = \vec{a} + \lambda(\vec{b} - \vec{a})$  vector form

$\frac{x-x_1}{x_2-x_1} = \frac{y-y_1}{y_2-y_1} = \frac{z-z_1}{z_2-z_1} = \lambda$  Cartesian form

Angle b/w 2 lines:  $\vec{r} = \vec{a} + \lambda \vec{b}, \vec{r} = \vec{c} + \mu \vec{d}$   
 $\cos \theta = \frac{|\vec{b} \cdot \vec{d}|}{|\vec{b}| |\vec{d}|}$

Shortest distance b/w 2 lines: [shortest dist = 0 if intersecting]  
 $\rightarrow$  if parallel  $\rightarrow \vec{r} = \vec{a} + \lambda \vec{b}, \vec{r} = \vec{c} + \mu \vec{d}$   
 $\rightarrow$  then shortest dist =  $\frac{|(\vec{a} - \vec{c}) \times \vec{b}|}{|\vec{b}|}$   
 $\rightarrow$  if skew  $\rightarrow \vec{r} = \vec{a} + \lambda \vec{b}, \vec{r} = \vec{c} + \mu \vec{d}$   
 Shortest dist =  $\frac{|(\vec{a} - \vec{c}) \cdot \vec{b} \times \vec{d}|}{|\vec{b} \times \vec{d}|}$

If two lines are intersector, then  $[(\vec{a} - \vec{c}) \cdot \vec{b} \times \vec{d}] = 0$

**Cartesian form:**

$\frac{x-x_1}{a_1} = \frac{y-y_1}{b_1} = \frac{z-z_1}{c_1} \& \frac{x-x_2}{a_2} = \frac{y-y_2}{b_2} = \frac{z-z_2}{c_2}$   
 Shortest dist =  $\begin{vmatrix} x_1 - x_2 & y_1 - y_2 & z_1 - z_2 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix}$   
 $\begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix}$

Plane passing thru a point  $\vec{a}$  & normal vector  $\vec{n}$ :



$(\vec{r} - \vec{a}) \cdot \vec{n} = 0$   
 Cartesian form:  $r = (x, y, z), a = (x_1, y_1, z_1), \vec{n} = (a, b, c)$   
 $a(x-x_1) + b(y-y_1) + c(z-z_1) = 0$


Plane passing thru 3 points  $\vec{a}, \vec{b}$  &  $\vec{c}$ :  $[\vec{AP} \ \vec{AB} \ \vec{AC}] = 0$

Plane passing thru points & parallel to vector  $\vec{b}$  &  $\vec{c}$ :  
 $[(\vec{r} - \vec{a}) \cdot \vec{b} \times \vec{c}] = 0$

Plane passing thru point & a line  $\vec{r} = \vec{c} + \lambda \vec{b}$ :  
 $[(\vec{r} - \vec{a}) \cdot (\vec{a} - \vec{c}) \times \vec{b}] = 0$




**Planes:**

Normal form:  $\vec{r} \cdot \hat{n} = d$




Cartesian form:  
 $lx + my + nz = 0$   
 $r = x, y, z$   
 $\hat{n} \rightarrow l, m, n$

Angle b/w two planes:



Foot of normal and image



$\frac{x_2 - x_1}{a} = \frac{y_2 - y_1}{b} = \frac{z_2 - z_1}{c} = - \frac{ax_1 + by_1 + cz_1 + d}{a^2 + b^2 + c^2}$   
 $\frac{x_3 - x_1}{a} = \frac{y_3 - y_1}{b} = \frac{z_3 - z_1}{c} = - \frac{2|ax_1 + by_1 + cz_1 + d|}{a^2 + b^2 + c^2}$

Convert the eq<sup>n</sup> of line  $(2x - y + z = 1 \text{ \& } x + y + 2z = 0)$  in Cartesian form.

$\rightarrow$  let  $z = t$   


$$\begin{aligned} 2x - y &= 1 - t \\ x + y &= -2t \\ \hline 3x &= 1 - 3t \\ x &= \frac{1 - 3t}{3} \end{aligned}$$

$$y = -\frac{1 - 3t}{3} \rightarrow t = \frac{1 - 3x}{3}$$


$$t = \frac{1 - 3y}{3}$$

$$\therefore x = \frac{1 - 3x}{3} = \frac{-1 - 3y}{3} = z$$

$$\Rightarrow \frac{x - \frac{1}{3}}{-1} = \frac{y + \frac{1}{3}}{-1} = \frac{z - 0}{1}$$



Angle b/w plane & line:



Sphere:  $x^2 + y^2 + z^2 + 2ux + 2vy + 2wz + d = 0$   
 centre  $(-u, -v, -w)$   
 radius  $= \sqrt{u^2 + v^2 + w^2 - d}$

$\rightarrow$  Diametric point:  
 $(x - x_1)(x - x_2) + (y - y_1)(y - y_2) + (z - z_1)(z - z_2) = 0$




# PROBABILITY

Probability of an event E,  $P(E) = \frac{\text{Favourable outcomes}}{\text{Total no of outcomes}}$

$0 \leq P(E) \leq 1$

$P(E) = 0 \rightarrow$  Impossible event


$P(E) = 1 \rightarrow$  Certain event



$P(A \cup B) = P(A) + P(B) - P(A \cap B)$   
 $A \cup B \rightarrow A \text{ or } B \quad | \quad A \cap B \rightarrow A \text{ and } B$

$P(E) + P(\bar{E}) = 1$

- Mutually exclusive Events: Only one of the events can occur at a time.  $P(A \cap B) = 0$
- Mutually independent Events: Occurrence of one event doesn't affect other events.  $P(A \cap B) = P(A) \times P(B)$
- Conditional Probability: Prob of A given that B has already occurred,  $P(A|B) = \frac{P(A \cap B)}{P(B)}$  / 4 suits  $\rightarrow$  Heart, Spade, Club, Diamond  
 Prob of B \_\_\_\_\_ A \_\_\_\_\_,  $P(B|A) = \frac{P(A \cap B)}{P(A)}$



• Baye's Theorem:

$$P(E/E_2) = \frac{P(E) \times P(E_2/E)}{P(E_1)P(E_2/E_1) + P(E)P(E_2/E)}$$

- E  $\rightarrow$  favoured event
- $E_1 \rightarrow$  opposite of favoured event
- $E_2 \rightarrow$  Already happened event.